

Predisposition in Plant Disease: Exploiting the Nexus in Abiotic and Biotic Stress Perception and Response

Richard M. Bostock, Matthew F. Pye,
and Tatiana V. Roubtsova

Department of Plant Pathology, University of California, Davis, California 95616;
email: rmbostock@ucdavis.edu, mfppe@ucdavis.edu, tvroubtsova@ucdavis.edu

Annu. Rev. Phytopathol. 2014. 52:517–49

First published online as a Review in Advance on
June 23, 2014

The *Annual Review of Phytopathology* is online at
phyto.annualreviews.org

This article's doi:
10.1146/annurev-phyto-081211-172902

Copyright © 2014 by Annual Reviews.
All rights reserved

Keywords

abscisic acid, cross talk, induced susceptibility, phytohormones, salinity

Abstract

Predisposition results from abiotic stresses occurring prior to infection that affect susceptibility of plants to disease. The environment is seldom optimal for plant growth, and even mild, episodic stresses can predispose plants to inoculum levels they would otherwise resist. Plant responses that are adaptive in the short term may conflict with those for resisting pathogens. Abiotic and biotic stress responses are coordinated by complex signaling networks involving phytohormones and reactive oxygen species (ROS). Abscisic acid (ABA) is a global regulator in stress response networks and an important phytohormone in plant-microbe interactions with systemic effects on resistance and susceptibility. However, extensive cross talk occurs among all the phytohormones during stress events, and the challenge is discerning those interactions that most influence disease. Identifying convergent points in the stress response circuitry is critically important in terms of understanding the fundamental biology that underscores the disease phenotype as well as translating research to improve stress tolerance and disease management in production systems.

INTRODUCTION

Abiotic stresses can dramatically alter the outcome of plant-pathogen interactions, and depending on the pathosystem and stress intensity, the stress may enhance or reduce disease. Recognition of the importance of predisposing environmental stress in plant pathology goes at least as far back as 1874, when Sorauer et al. formally introduced the concept and later discussed predisposition in their *Manual of Plant Diseases* (172). Hartig also recognized the importance that abiotic stress plays in altering the proneness of the host plant to disease, using the terms “predisposition” and “tendency to disease” interchangeably (73). Yarwood defined predisposition as “the tendency of nongenetic factors, acting prior to infection, to affect the susceptibility of plants to disease” (205). Implicit, and important, in the latter definition is that predisposing stresses can shift the outcome toward resistance or susceptibility.

The quest to improve plant tolerances to abiotic and biotic stresses is an ongoing effort. What is probably not fully appreciated, however, is that episodes of relatively mild abiotic stress can override disease resistance. Apparent from field experience and from the literature is that abiotic stress levels that are thresholds for predisposition occur routinely in agricultural, forest, and nursery production systems. Furthermore, in the absence of a pathogen, plants often recover rapidly and fully when the stress is relieved (27). This is significant for plant breeding and in food and fiber production in which abiotic stress can (*a*) reduce inoculum thresholds necessary for disease, (*b*) degrade the consistency and reliability of pathogenicity tests and disease resistance screens, and (*c*) diminish or nullify the efficacy of other disease management measures. Although the problem of predisposing stress in plant diseases is not new, the specter of climate change adds urgency for a better understanding of abiotic and biotic stress interactions (65).

SCOPE OF REVIEW

This review examines recent progress in our understanding of the interaction between abiotic stress and disease, and builds upon themes presented previously on cross talk and trade-offs in phytohormone signaling in relation to induced resistance (24). Here, we emphasize stresses that predispose plants to levels of pathogen inoculum that would not be damaging in the absence of the stress and discuss how phytohormone networks may engage unproductively to compromise the host plant. Although abiotic stresses affect pathogens, we do not delve into this aspect, except to mention it as an experimental consideration. For additional background, the reader is referred to excellent reviews and the references therein that address plant abiotic stress perception, signaling, and response (143, 208), and others that address biotic and abiotic stress interactions in plants (13, 148, 157). Desprez-Loustau et al. (45) thoroughly cover the predisposing effect of drought in diseases of forest trees. For population- and landscape-level perspectives and the impact of climate change on plant diseases, the reader is referred to an excellent special issue of the journal *Plant Pathology* (33).

ABIOTIC AND BIOTIC STRESSES AND THE DILEMMA OF SHARED SIGNAL-RESPONSE NETWORKS

Stress can be defined as “a sudden change in the environment that exceeds the organism’s optimum and causes homeostatic imbalance, which must be compensated for” (92). Stress imposes strains on a biological system that can be distinguished as plastic or elastic (15, 181). Strains that are plastic result in irreversible physical or chemical changes that often lead to death of the plant. Elastic strains impose physical or chemical changes that are reversible when the stress is removed. Elastic strains, such as water deficit or the hypoxia incurred by waterlogging, may have consequences that become irreversible with sufficient duration (15).

As sessile organisms, plants have effective physiological mechanisms to maintain homeostasis during stress events. These are adaptive, at least initially, in order to promote the plant's health and survival (96, 97). The dilemma that arises in predisposition, however, is that an adaptation in one context, such as adjustment to water deficit, may be maladaptive when there is a need to also resist pathogens and pests. The notion of a response hierarchy attempts to rationalize the apparent dominance of one response over another in the face of concurrent stresses (71). An evolutionary interpretation is that the dominant response has undergone stronger selection pressure, for example, a trait arising from the continual need for plants to minimize water stress or damage from UV irradiation versus a threat of a sporadic nature, as attack by pathogens might be viewed. The notion of accepting alternative evolutionary scenarios was raised previously in the context of why some current traits may not result in the highest fitness, the example being trade-offs we observe in defense signaling when plants are challenged by different attackers (24). Such traits may have been pulled along by pleiotropy, linkage with a correlated trait, genetic drift, and/or lack of variation. The degree to which the stress response circuitry is shared to counter an abiotic stress but not the biotic stress, and vice versa, could be a result of these genetic processes.

A general adaptation syndrome (GAS) posits a common stress response in plants to evoke similar coping mechanisms (104). Responses common to anoxia, drought, heat, chilling, flooding, salinity, desiccation, and freezing, which include both physiological and morphological adaptations, support this view (149). Numerous investigations, including contemporary transcriptome, proteome, and metabolome studies, reveal that abiotic and biotic stressors engage common signals and share responsive genes and products (34, 97, 113). Plant stress response pathways are conserved (161) and are viewed as highly integrated, overlapping, and nonlinear to balance and optimize plant performance in the face of diverse challenges (59). Different stressors may cause similar cellular damage to initiate the shared signal-response cascades through perturbation of ion channels and mechano- and osmosensors, membrane disruption, and generation of reactive oxygen species (ROS) (38, 141, 195) (**Figure 1**). Implicit and perhaps desired is that subfunctionalization downstream from the common signaling events will afford opportunities for tapping selective responses to improve tolerance to different stresses.

Certainly, positive interactions and synergies occur, examples being cross tolerance to chilling following adaptation to water stress and a mild stress that may enhance disease resistance (176). It is important to acknowledge the positive effects of environmental stresses, particularly when the stress is imposed gradually rather than abruptly, thereby enabling the physiological adjustments necessary for tolerance (96). The GAS model is instructive in that the stress response is viewed in three stages: an initial alarm reaction, an acclimation stage during which a degree of tolerance is acquired, and, if the stress persists, an exhaustion phase resulting in collapse or severe compromise (**Figure 2**).

EXAMPLES OF PREDISPOSING ABIOTIC STRESSES

Plant pathology textbooks include a section on abiotic diseases with discussion of water stress, heat, chilling, freezing, light quality, air pollution, nutrient deficiency or excess, salinity, and herbicide injury as well as other stresses. Any of these at sufficient intensity can damage or kill plants, and this damage can be further exacerbated by pathogen attack. All of these at intensities below an irreversible damage threshold, and often in the absence of visible injury, can predispose plants to pathogens. This is to say that predisposition occurs with both plastic and elastic strains. However, of special interest are mild elastic strains that occur with brief or episodic stress from which the plant normally recovers (**Figure 2**). Such reversible strains present experimentally tractable




Figure 1

General signal-response sequence in plants following abiotic and biotic stress events that illustrates the concept of shared pathways and phytohormone cross talk to shape transcriptional reprogramming and response. Factors involved in signal-response coupling include various kinases, phosphatases, transcriptional activators and repressors, Ca^{2+} signaling elements, and so forth (*light orange box*), and downstream outputs include gene expression and biochemical and physiological responses (*light blue box*). Dashed arrows indicate feedback, which could be positive or negative, to enhance or attenuate reactive oxygen species (ROS) and phytohormone action. Adapted from Fujita et al. (62). Abbreviation: ABA, abscisic acid.




Figure 2

Model of the plant response to abiotic and biotic stresses that integrates features of Selye's general adaptation syndrome, concepts of strain severity and duration, and disease predisposition. The phase of greatest vulnerability to disease occurs immediately following or soon after a stress event and prior to onset of acclimation. The dashed line illustrates predisposition and abrogation of stress adaptation as disease develops. In the absence of the abiotic stress, the plant would normally resist the pathogen or at least the level of inoculum present. Adapted from Leshem et al. (104).

approaches for mechanistic inquiry and, relative to studies of severe stress, may afford greater opportunities to identify genetic strategies to mitigate predisposition.

A classic review in this journal (165) provided a generalized model for predisposing stress that accommodated different outcomes, either enhanced resistance or susceptibility, depending on the parasitic strategy of the pathogen. The underlying paradigm interprets these different outcomes as a gain or loss in resistance resulting from the stress's impact on host defenses. However, it is fair to note that the stress may also cause a gain or loss in susceptibility, attributable to what might be viewed as susceptibility factors, such as cell death programs (49). The difference may seem nuanced and semantic, but the paradigm selected can influence the framing of hypotheses and the selection of analytical targets.

Abiotic stress generally results in reduced severity or incidence of diseases caused by obligate, or biotrophic, pathogens (3, 166), although there are exceptions, such as diseases caused by some viruses, fungi, and nematodes (177, 180). Abiotic stresses can predispose plants to potentially aggressive hemibiotrophic pathogens, resulting in severe disease from very low levels of inoculum. Notable here are *Phytophthora* root and crown rots, where episodes of plant water stress in its various forms can be a critical determinant for disease development and severity (56, 99, 115) (**Figure 3**). Perhaps the most pronounced impact of abiotic stress is to facilitate diseases caused by weakly aggressive facultative pathogens and those usually present in association with their hosts as saprophytes or endophytes (45). These include the root- and crown-infecting pathogens *Pythium ultimum* and *Fusarium* spp.; pathogens of aerial parts, such as *Alternaria* spp. and *Botrytis cinerea*; and many canker-causing pathogens of trees and woody perennials (45, 118, 177) (**Figure 4**).

An analysis of recent literature by Cramer et al. (40) found that of the more than 35,000 papers published between 2001 and 2011 on abiotic stress, 14% dealt with water stress, 28% with temperature stress, 22% with light stress, and 35% with chemical/soil stresses (e.g., nutrients and minerals, salinity, air pollutants). In terms of their importance as factors in disease predisposition, water, nutrient/salinity, temperature, and air pollution, stresses are perhaps the most common environmental triggers. Variation in light intensity or duration is generally not of significance to impact diseases in the field but can be a consideration in highly managed production or experimental systems, in natural ecosystems in studies of understory plants, or in combination with other stresses (58, 165). Subtoxic levels of synthetic herbicides can also increase or decrease disease in crops in the field, particularly those caused by facultative pathogens, and are commonly reported (55). Direct effects of the herbicide on the pathogen have also been reported.

Physiological and Experimental Considerations

As with any experimental pathosystem developed for mechanistic inquiry, the environmental conditions and disease assays in a predisposition study must be carefully considered and optimized. Typically, one environmental parameter is isolated and varied, and disturbing influences that may confound interpretation are eliminated or minimized. To discern stress treatment effects on the plant-pathogen interaction, it is desirable to use plants at the same developmental stage with uniform growth conditions and maintain sampling consistency while using a sensitive and quantitative disease assay. Diurnal changes and circadian phases are also important to consider, as these influence gene expression and response during stress events (63, 198). Whether plants are actively transpiring can also influence the impact of a stress on host physiology, an example being salinity (an accumulation of salts in leaves is higher as a result of transpiration) (134). Most contemporary studies incorporate biochemical and molecular methods that can be exquisitely sensitive to subtle external changes, so experiments are typically conducted within tightly




Figure 3

Predisposition in *Phytophthora* root and crown rot. (a) Impact of salinity stress prior to inoculation in four *Phytophthora*-plant interactions: *Phytophthora capsici*-tomato (*Solanum lycopersicum*); *Phytophthora cryptogea*-*Chrysanthemum*; *Phytophthora ranorum*-*Rhododendron*; *P. capsici*-*Arabidopsis*. Roots were exposed to 0.2 M NaCl/0.02 M CaCl₂ in half-strength Hoagland for 16–24 h, rinsed and returned to half-strength Hoagland to recover, and then inoculated with zoospores (10^4 – 10^5 per ml). Controls included no salt and noninoculated treatments as indicated. Images taken after an appropriate incubation period. Portions of panel a adapted from References 50 and 159. (b) Impact of an episode of heat stress on development of *Phytophthora* root rot caused by *P. cryptogea* in potted *Chrysanthemum* plants. Potted plants were removed from the greenhouse and placed in an outdoor nursery for one day either in partial shade or in full sun, achieving maximum soil temperatures as indicated. Plants were then inoculated with a zoospore suspension and returned to the greenhouse; several days later the roots were washed for isolations and examined for symptoms. Note the discoloration and necrosis of the inoculated roots from the pots exposed to full sun. Images in panel b courtesy of J.D. MacDonald (117).




Figure 4

Examples of predisposition to fungal canker diseases in trees. (a) Dieback of raywood ash (*Fraxinus oxycarpa* Raywood) following drought stress caused by *Botryosphaeria stevensii*. (b) *Botryosphaeria* canker development in inoculated nonstressed (stem water potential = -1.0 ± 0.5 MPa) and water-stressed (-3.1 ± 0.4 MPa) raywood ash branch segments. Arrows indicate canker lengths. Photos in panels a and b courtesy of T.R. Gordon. (c) Almond (*Prunus dulcis*) stems from cold-stored seedling trees showing symptoms and signs of cankers caused by *Fusarium* spp. and *Cylindrocarpon* spp. (d) Influence of bark moisture measured as relative bark turgidity on the susceptibility of dormant almond tree segments to *Fusarium acuminatum*. Branch segments were inoculated following a 14-day desiccation period (84% relative bark turgidity) or without desiccation (88% relative bark turgidity) and then examined 14 days after inoculation. W,C: wound-only control; W,I: wound inoculation. Panels c and d adapted with permission from Reference 118.

controlled growth environments. We refer the reader to excellent reviews of experimental design and analytical considerations for studies of water stress (189) and stress transcriptomes (92).

Adjusting the inoculum to an appropriate titer is also an important consideration. Inoculum density that is too high can mask the predisposing effect of a stress, so levels are adjusted to where there is little or no disease in the absence of the stress. Plants may resist relatively high inoculum levels, as with weak, facultative pathogens, and it is only with predisposing stress that disease manifests. This can be an important consideration for establishing etiology, whereby Koch's postulates should be performed, if necessary, on both nonstressed and moderately stressed

plants to evaluate the potential of a species to cause disease in a given host (165). Predisposing stresses can also affect pathogen physiology or behavior (57), and, if so, this needs to be considered in the experimental design. The pathogen may be physiologically compromised by the imposed stress, inactivating inoculum or rendering it temporarily incompetent until favorable conditions are restored. The pathogen may be stimulated by the stress, as with salinity (115) and flooding (99) to increase colonization of roots by *Phytophthora* spp. Isolating the stress effect to the plant can be done experimentally to some degree by separating in time or place the stress treatment from the inoculation (50, 152). The time course over which stress is imposed can influence how the plant responds. Where plants are grown in containers, roots proliferate at the sides and bottom of the pot. When water is withheld it disappears quickly from these locations so stress comes on much more quickly than it would under field conditions. Consequently, plants cannot adjust physiologically to the same degree, and the effects of water stress may overwhelm subtler effects on disease susceptibility than might otherwise be apparent.

Water Stress in Predisposition

Abiotic stresses that impact plant water potential (ψ_w) have provided an important focus in predisposition research (27) and include dehydration from water deficit, hypoxia from waterlogging, and osmotic stress from soil salinity. Chilling and freezing, in addition to their direct injury to plant membranes and other structures, also impact plant ψ_w (181). Classic studies of fungal canker diseases of trees (165) demonstrated predisposition when bark moisture dipped below a critical threshold of approximately 80% relative turgidity, although higher bark turgidities (e.g., 84% in almond stems) can be predisposing to fungal pathogens (118) (**Figure 4**). In general, stem xylem potentials of -1.2 to -1.5 MPa provide a threshold for predisposition to nonaggressive canker fungi during episodes of water stress in woody plants (167). Desiccation of woody tissues due to drought stress leads to loss in the mechanical strength of the bark-wood bond (131) and can result in bark cracks that can be invaded by opportunistic endophytes (21). Transplanted tree seedlings can also experience a severe physiological shock due to injury to root systems and consequent loss in capacity for water absorption (96), resulting in the activation of cryptic infections (118).

Root and crown diseases caused by *Phytophthora* spp. and other soilborne pathogens have provided models whereby cycles of soil saturation, drought, and salinity were shown to be dominant predisposing factors to markedly increase disease severity (39, 42, 56, 115). This research defined many of the physiological parameters of stress-induced predisposition in the laboratory, greenhouse, and field, and demonstrated that predisposition can occur as a result of relatively minor stress. Brief exposure to water potential deficits from -0.5 to -1 MPa predisposes various plant species to *Phytophthora* root and crown rots (**Figure 3**). Waterlogging compromises the plant, although the impact varies across species (30, 168). Tomato and tobacco plants will show signs of wilting and leaf yellowing within a few hours of flooding, indicative of the rapid dehydrating effect and impact on hydraulic conductance (81). These studies consider stress effects on both the host and the pathogen and have shown that the *Phytophthora* spp. of concern function effectively in soil during (or, in the case of drought, immediately after) the stresses that cause predisposition. This research has raised grower awareness of the need to optimize watering regimes where possible and brought attention to the importance of predisposing stress in assessing plant performance against certain pathogens, resulting in the incorporation of stress-screens in disease resistance breeding programs (76).

For typical mesophytic crop plants, as the ψ_w declines over the range -0.2 to -2.0 MPa, which encompasses plants that are well-watered to plants experiencing mild water stress, cell expansion




Figure 5

Predisposition thresholds in relation to water potential and major physiological and biochemical changes in plants under water stress. Cell expansion and wall and protein synthesis are most sensitive to a decline in water potential, followed by stomatal conductance and photosynthesis. Abscisic acid (ABA) and osmocompatible solutes begin to accumulate at the threshold for predisposition. Adapted with permission from References 167 and 181.

and wall and protein synthesis are most sensitive, declining rapidly within this range (**Figure 5**). This rapid decline in protein synthesis (48) would seem to compromise the defense budget of the plant and is probably underappreciated in contemporary studies that focus on the dramatic redirection of gene expression and compensatory metabolic adjustments during the stress response. Photosynthesis and stomatal conductance subsequently decline with concomitant accumulation of solutes and induction of the phytohormone abscisic acid (ABA) (181). In many plants, root:shoot ratios increase as part of the strategy to adapt to reduced water availability.

Temperature and Other Factors in Predisposition

Temperature extremes, similar to severe water stress, can create a plastic strain that is directly injurious, particularly when they occur abruptly and the plant is insufficiently acclimated. Such injuries provide infection courts for facultative pathogens. Generally, freezing temperatures between -20°C and -30°C can predispose woody plants in temperate climates to fungal canker diseases and diebacks (165). Mild to moderate freezing contributes to development of bacterial canker and short-life disorders in stone fruit trees (29).

Temperature stresses of a milder form provoke elastic strains that increase disease proneness. Chilling injury can occur in plants at temperatures $>0^{\circ}\text{C}$ and up to 15°C . Chilling is a key predisposing factor in cotton leaf senescence and susceptibility to *Alternaria alternata*, leading to premature defoliation (210), and tropical and subtropical plants are typically quite sensitive to chilling. In the rice blast pathosystem, cold temperature and ABA suppress resistance to

Magnaporthe oryzae (94). Cold temperatures induce water stress, wilting and leaf necrosis, and ABA (184). Nonetheless, transcriptome analysis shows both positive and negative regulation of gene expression by ABA in chilled pepper plants, and ABA application can alleviate some symptoms of chilling injury (69). The physiological changes occurring during cold acclimation can also enhance disease resistance in some plants, such as occurs in various winter cereals to snow mold diseases (101). Salicylic acid (SA) biosynthesis is induced by cold temperatures with a corresponding induction of pathogenesis-related (PR) proteins, possibly as a preemptive strategy to minimize the impact of potential infections of injured host tissue (93).

The integration of downstream signaling generated by the seemingly disparate stresses of cold and pathogen infection is evident from studies with a chilling sensitive mutant (*chs2*) in *Arabidopsis* in which a shift from 22°C to temperatures below 16°C causes a hypersensitive response (HR)-like cell death accompanied by electrolyte leakage, PR gene expression, and ROS and SA accumulation (79). *CHS2* encodes the R protein RPP4, which is effective against the downy mildew pathogen *Hyaloperonospora arabidopsidis*. A single amino acid substitution in the nucleotide binding domain confers the temperature-dependent deregulation of RPP4 in the *chs2* mutant that results in chilling sensitivity and the associated reactions that mimic a response to infection. The integration of cold and canonical defense responses is further evidenced by shared regulators and transcriptional repressors and activators (187).

Transient episodes of heat stress can increase susceptibility or resistance to pathogens. In nurseries, pots exposed to direct sunlight can achieve soil temperatures of 45°C or more, which can cause heat injury to ornamental roots and significantly increase severity of *Phytophthora* root rot (117) (**Figure 4b**). Even mild temperature elevation can nullify host resistance, the classic examples being interactions conditioned by temperature-sensitive *R* genes (212). Elevated temperatures can also induce resistance, as in cucumber seedlings exposed to a heat shock of 50°C for 40 s (174). Heat-shock factors, which recognize consensus sequences in the promoters of heat-induced genes, and PR and heat-shock proteins are associated with heat shock-induced resistance in various species (100).

Although all mineral nutrients in inappropriate amounts can impact severity of disease (53), the impact of nitrogen (N) fertility is the most commonly reported and studied (80). Although excess N often increases disease, insufficient N can have a similar effect (171). Also important is the form of N as ammonium or nitrate because each can affect diseases differently. In addition to N form, a number of factors contribute to N's impact, including genetics of the host, applied N rate and availability, and the influence of soil pH and redox on the availability of other ions (80). Ammonium-N increases *Fusarium* wilt of tomato, particularly in acidic soils, and this can be managed with lime and nitrate-N fertilizer, a treatment that also decreases the availability of Mn and Fe. However, wheat take-all disease is reduced by ammonium-N, acidic soil, and Mn. N stress, whether resulting from insufficient or excessive N, results in a decline in photosynthesis and reduced growth (35). N deficiency consistently results in increased ABA levels in plants (35), as does excessive N in the form of ammonium-N (105). There is also a range within which increased N enhances growth and disease susceptibility, as in stem rot of rice (90) and pitch canker of Monterey pine (109). This is interpreted as resulting from a trade-off between growth and defense, which includes less lignification and greater succulence of the tissue.

Atmospheric pollutants, notably SO₂, NO_x, and ozone (O₃), are directly injurious to plants but also indirectly influence plant-pathogen interactions (19). O₃, for example, can increase severity of diseases caused by necrotrophic pathogens and reduce severity of those caused by biotrophs, although this effect can be quite variable (162). The physiological and cellular impacts of O₃ in plants are similar to the effects of treatment with elicitors, with the associated induction of antioxidant systems, expression of PR proteins, and other responses.

PHYTOHORMONE NETWORKING IN PREDISPOSITION

Phytohormones, together with ROS, provide important signals to help orchestrate the similar signaling cascades, transcriptional changes, and metabolic and cellular responses to abiotic and biotic stresses (24, 95, 157). Stress-induced changes in both concentration and perception of phytohormones are important to consider in conducting and interpreting these studies. ABA, jasmonic acid (JA), ethylene (ET), SA, auxin [indole-3-acetic acid (IAA)], and cytokinins (CKs) figure prominently in the literature on disease resistance and susceptibility (157). More recently, gibberellic acid (GA) and its interaction with DELLA proteins as well as the brassinosteroids (BRs) have gained interest in studies of abiotic-biotic stress interactions (18, 137, 146). Whereas SA, JA, and ET have shaped much of the current understanding of induced resistance, it has become evident that there is considerable cross talk among all the major phytohormones in abiotic and biotic stress signaling and response. Phytohormone and ROS modulation of immunity in plant-microbe interactions have been reviewed recently (126, 148, 157, 191). These excellent reviews cover much of the territory, so here we discuss the aspects most germane to predisposition toward susceptibility, with an emphasis on ABA and its interactions (**Figure 6**).

Abscisic Acid


ABA is a highly conserved stress-related signal that occurs throughout all kingdoms except Archaea and appears to be universally triggered under conditions of limited cellular water availability (74). In higher plants, ABA, a C₁₅ isoprenoid, is synthesized by the plastidial 2-C-methyl-D-erythritol-4-phosphate (MEP) pathway via cleavage of C₄₀ carotenoids (136) (**Figure 7**). Discovery and characterization of the PYR/PYL/RCAR family of ABA receptors and the core signaling pathway are leading to a deeper understanding of ABA action within different developmental and environmental contexts (41) and revealing opportunities that may lead to improved stress tolerances in plants (83, 202). Many stress responses engage ABA, and, remarkably, it is estimated that as many as 10% of all protein coding genes in *Arabidopsis* show differential expression following ABA treatment. Many of these genes respond to abiotic and biotic stresses (92). Several studies, including a meta-analysis of a series of stress transcriptomes (4), provide evidence for ABA as a signal for defense-related gene expression (119). ABA-responsive elements are highly represented in the promoters of many of these genes, although ABA alone may be insufficient to fully engage their expression and, in some cases, may even suppress expression.

ABA's primary role in stress responses is to evoke adaptive physiological changes toward water balance and cellular dehydration tolerance (38, 74, 188), which includes guard cell regulation (86). ABA, acting in concert with other phytohormones, helps maintain root growth while reducing shoot growth and photosynthetic rate during stress episodes, alters the capacity for nutrient uptake, and invokes gene expression leading to protective proteins (e.g., LEA proteins and dehydrins) and osmocompatible solutes (35, 78).

Abscisic Acid in Disease

A causal role for ABA in disease is supported by studies with ABA-modified plants (12, 119, 162). Early investigations of ABA action in plant-microbe interactions in potato (75) and soybean (31, 120, 127) demonstrated that ABA pretreatment abrogated race-cultivar resistance to *Phytophthora* spp. and suppressed phytoalexin accumulation and defense responses, with ABA-synthesis inhibitors enhancing resistance. ABA predisposes wheat seedlings to *Fusarium culmorum*, causal agent of foot rot, a classic drought-incited disease (123), and pine seedlings to *Cylindrocarpon destructans*

(98). Most studies now incorporate ABA-synthesis and -perception mutants, and some have examined the connection between abiotic stress, ABA, and disease (50, 94, 183, 206). In *Arabidopsis*, ABA treatment and drought stress induced susceptibility to *Pseudomonas syringae* pv. *tomato* (*Pst* DC3000) (128). The ABA-deficient *aba1-1* mutant displayed reduced susceptibility to compatible isolates of the downy mildew pathogen *H. arabidopsidis*. An ABA-insensitive mutant, *abi1-1*, which has wild-type levels of endogenous ABA, was not altered in its host reaction to *Pst* or to *H. arabidopsidis*, although it is possible that other *ABI* genes may have compensated (41). There is also strong evidence that *Pst* DC3000 co-opts ABA biosynthesis and signaling in *Arabidopsis*, apparently through the action of its AvrPtoB effector, to promote disease in compatible interactions



(47). Whether targeting of host ABA signaling is a strategy used more widely among diverse pathogens remains to be determined.

Roots most directly encounter the predisposing impacts of waterlogging, salinity, and drought. Studies of root-pathogen interactions suggest shared elements and common themes in stress and defense network dynamics with leaves, even if details differ qualitatively and quantitatively. An elegant study of cell type-specific transcriptional responses in *Arabidopsis* roots found that ABA marker genes, unlike those for other phytohormones, are induced by salt stress in a semiubiquitous manner in all cell layers of the root (51). Because of the importance of predisposing stress in soilborne diseases, we have examined ABA-regulated susceptibility in roots with experimental treatments that are informed by predisposition in the field (23, 56, 116).

The tomato (*Solanum lycopersicum*)–*Phytophthora capsici* interaction provides an important experimentally tractable crop model that can give a clear disease phenotype within 48–72 hours post infection in hydroponic formats. Seedlings are predisposed by brief root immersion in saline solutions, followed by rinse and then return to a standard hydroponic solution and inoculation with zoospores (50). Disease and host colonization are evaluated by seedling collapse and qPCR of pathogen DNA. The acute stress regime results in a rapid phase 1 response as defined in salinity research (133), and noninoculated but stressed plants recover completely. In tomato roots, ABA increases rapidly, peaking 6 to 12 hours after salt treatment, and then declines (50), although the timing may vary depending on the size and age of the plant. ABA induction precedes or temporally parallels the onset of the predisposed state, which is evident within 4–6 hours of salt exposure and persists for up to 24 h after salt removal, well after the decline in ABA levels to near prestress levels. Salinity stress in roots and its systemic impact are evident in various interactions, with similar effect in leaves challenged with the bacterial speck pathogen *Pst* (152, 183) and in soilborne infections by *Phytophthora* spp. in several ornamental species (50, 159).

Roots and shoots of wilted ABA-deficient tomato mutants accumulate only a fraction of the ABA present in wild-type plants and do not show the adaptive responses to water stress. Following a

Figure 6

Phytohormone interactions in abiotic and biotic stress signaling and response. In each panel, black arrows indicate a positive or feed-forward effect and red blocked lines indicate inhibition. Dashed lines imply indirect or multiple steps in the sequence. Factors involved in signal-response coupling (e.g., kinases, phosphatases, transcription factors, etc.) are in light orange boxes, and downstream outputs (gene expression and responses) are in light blue boxes. The models are not intended to be comprehensive and are derived from those presented in detail in References 148, 153, 157, 196, and 209. (a) Salicylic acid (SA) path for induced resistance and antagonism between SA and abscisic acid (ABA). SA has various effects on abiotic stress tolerance as well. (b) Ethylene (ET) and jasmonic acid (JA) and the ERF (ethylene responsive factor) and MYC response branches. (c) Cytokinins (CKs) counteract ABA responses to water stress, but CKs can also enhance stress tolerance. In *Arabidopsis*, CK signal perception and transduction are mediated by a two-component histidine-kinase phosphorelay system composed of *Arabidopsis* histidine kinases (AHKs), histidine phosphotransfer proteins (AHPs), and response regulators (ARRs), with members of the latter providing positive and negative feedback regulation of CK signaling. (d) DELLAs act as repressors of gibberellic acid (GA) signaling and of JAZ (JASMONATE-ZIM DOMAIN) repressors, the latter resulting in the release of JA-mediated gene expression and cross talk with SA. (e) Auxins, such as indole-acetic acid (IAA), as well as ABA, SA, and biotic and abiotic stressors, can induce the GH3 auxin-amino acid-conjugating enzyme WES1 to lower free auxin pools in the cell. SA can also interfere with auxin-mediated gene expression. (f) Brassinosteroids (BRs) can mitigate abiotic stresses. They share BAK1 (BRI-associated receptor kinase) with the PAMP (pathogen-associated molecular pattern) flg22, and thus may compete for BAK1's attention. BAK1 complexes with other receptor-like kinases (RLKs), leading to downstream effects. Abbreviations: ABI5, abscisic acid insensitive 5; ANAC, abscisic acid-responsive NAC; ARF, auxin response factor; Asp, aspartic acid; BRI1, brassinosteroid insensitive 1; DELLA, DELLA protein; GH3, Gretchen Hagen 3; LOX2, lipoxygenase 2; MPK, MAP kinase; MYC, MYC transcription factor; NPR1, nonexpressor of pathogenesis-related genes; ORA, octadecanoid-responsive *Arabidopsis* AP2/ERF; PDF1.2, plant defensin 1.2; PR, pathogenesis related; RGL2, RGA-like 2 transcriptional regulator; SCF, Skp1-Cullin1-F-box; TGA, TGA transcription factor; TIR, transport inhibitor response; TRX, thioredoxin; VSPI, vegetative storage protein 1; WES1, Weso 1; WRKY, WRKY transcription factor.




Figure 7

Abscisic acid (ABA) biosynthesis and features of ABA signaling. Black arrows indicate a positive or feed-forward effect, and red blocked lines indicate inhibition. ABA biosynthesis and catabolism abbreviations: 2-MEP, 2-C-methyl-D-4-erythritol phosphate; ZEP, zeaxanthin epoxidase; NCED, 9-*cis*-epoxycarotenoid dehydrogenase; ABA2, short chain alcohol dehydrogenase; AAO, abscisic aldehyde oxidase; CYP707A, a family of ABA-8'-hydroxylases; AOG, ABA O-glucosyl transferases. Various biosynthesis mutants in *Arabidopsis* and tomato are indicated. ABA receptor-mediated signaling: binding of ABA to the PYR/PYL/RCAR family of receptors inhibits PP2Cs, a family of serine-threonine phosphatases (e.g., ABI1, ABI2, HAB1). PP2Cs normally keep SnRKs (SNF1-related protein kinases) at bay. In the presence of ABA, SnRKs phosphorylate transcription factors [ABFs (ABA response-element binding factors); e.g., ABI3, ABI5] to activate gene expression. Ubiquitin-proteasome system: selected E3 ubiquitin ligases that target various elements to promote (+) or inhibit (-) ABA signaling. PP2A is another family of serine-threonine protein phosphatases. See References 41, 54, 61, 111, and 136 for further details. Abbreviations: ABI, abscisic acid insensitive; AIP2, ABI3-interacting protein 2; AFP, ABI5 binding protein; ARIA, arm protein repeat interacting with AFB2; CHIP, C-terminal HSP interacting protein; KEG, keep on going E3 ligase; SDIR1, salt- and drought-induced RING finger 1; XERICO, a RING-H2 zinc finger protein, from the Greek, meaning "drought tolerant."

stress episode, these mutants show a significantly diminished predisposition to *P. capsici* relative to wild-type plants (50). The difference is most dramatic in the *sitiens* mutant, having the most depressed phenotype, with less than 10% of the ABA levels of wild-type seedlings. However, exogenous ABA only partially complements the *sitiens* mutant to full susceptibility (50), suggesting additional ABA-independent factors contribute to the predisposing effect of salt or, perhaps, synergy between salt and ABA in conditioning the plant response, as observed, for example, in rice cells (25). In fungal (11) and bacterial (10) diseases of tomato in which ABA pushes the interaction toward susceptibility, *sitiens* displays a rapid induction of defense responses that are otherwise diminished in the wild type.

Although there are many examples of ABA shifting the host-pathogen dynamic toward susceptibility (12), there are also examples in which abiotic stress and ABA enhance disease resistance (32, 185). Notable among these are powdery mildew (3) and other fungal diseases and several bacterial diseases. In the latter case, stomatal closure triggered by ABA in concert with SA may prohibit ingress of the pathogen (121), although recent evidence indicates an important role for ABA-independent oxylipin signaling in stomatal function during drought stress and pathogen infection (130, 163). Recognition of this complexity prompted a model that considers parasitic strategy (66), the timing and intensity of ABA signaling during the course of infection, and ABA's interaction with SA and JA to explain different outcomes (185).

Defense Network Signaling During Predisposing Stress: Does Abscisic Acid Trump Salicylic Acid, Jasmonic Acid, and Ethylene?

Changes in SA, JA, and ET that strongly modulate defense against biotic attackers coincide with changes in ABA during abiotic stress events (24, 181). Central to the downstream signaling that couples SA to induced resistance are the thioredoxins TRX-H3 and TRX-H5 and the transcription cofactor NPR1 (nonexpressor of PR genes 1). Monomerization and nuclear localization enable NPR1 to combine with members of the TGA subclass of bZIP family transcription factors (TFs) to activate defense gene expression (148, 191). However, SA has additional, and sometimes conflicting, physiological effects other than its importance in defense signaling (191). SA can increase or decrease plant tolerance to various abiotic stresses, depending on how much and where it is applied, the developmental stage of the plant, and the plant's overall redox status and stress history (78). SA can interfere with or synergize ABA- and osmotic stress-regulated gene expression and response (190, 206), disrupt the plant's ability to maintain water balance (17), induce partial tolerance to predisposing levels of salt stress (175), and contribute to redox homeostasis and protection against oxidative stress in various contexts (190) (**Figure 6a**).

ABA inhibits SA-mediated acquired resistance to *Pst* in *Arabidopsis* and tomato, whether the ABA is applied exogenously or induced by drought or NaCl stress (46, 60, 129, 183). Details of the cross talk between ABA and SA have been further delineated in *Arabidopsis* in a compelling study by Yasuda et al. (206), who used various signaling mutants and two resistance-inducing SA mimics, one that acts upstream of SA to stimulate its synthesis, and one that acts downstream of SA to engage its targets. ABA and NaCl stress inhibited the action of both SA mimics, with evidence from various mutants for reciprocal inhibition by SA of stress-induced ABA synthesis and ABA-regulated gene expression. The results implicate multiple points of interaction between ABA and SA as well as both dependence and independence of NPR1 in the outcome of the cross talk. ABA's inhibition of SA-mediated defenses also occurs in monocots, as recently reported in the interaction of rice with the bacterial leaf blight pathogen *Xanthomonas oryzae* pv. *oryzae* (201).

We examined the influence of SA on NaCl predisposition to *Pst* and *P. capsici* in wild-type and *nabG* transformed tomato seedlings that are compromised in SA-mediated resistance. In

both diseases, *nabG* plants were more susceptible than the wild type, with salt stress causing the expected additional increase in disease severity in both host backgrounds (152). However, the impact of predisposing stress in the *nabG* lines was proportionally the same as in the wild type. One interpretation is that the impairment of SA action following osmotic stress contributes little relative to the dominating impact of ABA and brings into question any direct role of SA-mediated processes to counteract predisposition within our brief stress regime.

JA and ET also interact with ABA and SA, although whether these interactions are positive or negative in target gene expression and response depends on the context (4, 7, 28, 44, 60). Various abiotic and biotic stresses induce JA and/or alter JA sensitivity to induce defense gene expression and adaptive responses. JA is an oxylipin derived from α -linolenic acid in plastids, with further metabolism in peroxisomes (197). JA is converted to its isoleucine conjugate, JA-Ile, which binds to the F-box factor COI1 in association with an SCF E3 ubiquitin ligase. This activated SCF^{COI1} complex then binds to JAZ (JASMONATE-ZIM DOMAIN) transcriptional repressors to release and target them for degradation by the proteasome, alleviating the block on transcription (197).

There appear to be two pathways through which JA acts in stress responses, the ERF (ethylene responsive factor) branch and the MYC branch, each with different outputs and regulation, that are mutually antagonistic (148) and are conserved in *Arabidopsis* and tomato (26) (**Figure 6b**). In the ERF branch, ET conspires with JA to induce expression of TFs, ERF1A/ORF59, and their target PR genes (e.g., *PDF1.2*), responses typically associated with resistance to necrotrophs. In the MYC branch, MYC TFs are induced by the coordinated action of ABA and JA to induce other TFs (ANAC019, ANAC055, and ANAC072) that engage gene expression to suppress SA accumulation, coordinate responses to water stress, and counter insect herbivory (89). MYC2 is induced by JA-JAZ derepression and serves as a core master regulator that forms homo- and heterodimers with MYC3 and MYC4 in complex with other proteins to orchestrate transcription in diverse JA responses (88). MYC2 also shows regulation by ABA and is required for ABA-dependent gene expression. The synergy of JA and ABA as translated through MYC2 may explain the enhanced disease resistance in some interactions.

The model that emerges is that JA-mediated stress responses are highly coordinated and finely tuned to address various challenges. There are many examples in which JA synthesis and signaling mutants display reduced resistance to bacterial, fungal, and oomycete pathogens and insect herbivores, affirming JA's position as a defense mediator (197). NPR1 is important in the negative cross talk between JA and SA (148), and a number of WRKY TFs, notably WRKY70, further contribute to modulating the SA-JA interaction to balance these different defense pathways (106). Although sophisticated circuitry engages JA to address different stresses, as with SA, it is unclear if JA and SA coordinate to counteract predisposition and how they negotiate ABA's strong induction during stress episodes. Also unresolved is the degree to which the antagonism between JA and SA is expressed to increase vulnerability to pathogens during stress events (24).

Molecular mechanisms governing induction and perception of ET as well as ET action and cross talk with other hormones in stress responses have been extensively studied (95). In *Arabidopsis*, stress-responsive mitogen-activated protein kinases (MAPK; i.e., MPK3 and MPK6) are important for inducing ET and phosphorylating specific ERF and WRKY TFs to trigger gene expression during infection (122). One of these targets, WRKY33, has been shown to be important for integrating collaborative signaling between JA and ET (22). ET can also be an important player in defense response activation by altering the JA-SA conflict. Depending on the timing of the ET signal during attack, NPR1's intermediary role can be circumvented (103) or SA's suppression of JA signaling can be abolished (102). Although ET interacts with multiple hormones and can modify disease resistance in plant-microbe interactions, including potentiating programmed cell death (PCD) (132), its contribution to predisposition is unclear. In tomato, we found that ET

was not induced by the brief episode of salt stress in our predisposition regime. Furthermore, the disease phenotype to *P. capsici* in the ET-insensitive *Never-ripe* mutant was not different from that in the wild type (50). Although ET can be induced in tomato during salt stress (6), this occurs well after ABA induction and far later than the time window of our stress regimes and disease assays. ET could be a factor during stress events of longer duration, and as with ABA and other hormones, ET is known to contribute to salt-stress adaptation (114), but additional research is needed to address any contributing role in predisposition.

Cytokinins Temper Abiotic Stress

CKs generally oppose ABA action in plants under water stress, with extensive cross talk between CKs and ABA and impacts on both ABA-dependent and ABA-independent stress responses (70). Short-term stresses, such as the acute salinity regime in our predisposition format, briefly elevate CK levels, but longer or more severe stress episodes reduce growth as well as CK levels and shift resources toward defense (**Figure 6c**). CKs are adenine derivatives containing either isoprenoid or aromatic side chains (112). The principal isoprenoid CKs are isopentenyl, *trans*-zeatin, *cis*-zeatin, and dihydrozeatin derivatives, and the rate-limiting enzymatic step in their synthesis is isopentenyl transferase (IPT). CK-modified tobacco plants in which IPT is transgenically expressed under control of the maturation- and stress-inducible promoter for the senescence-associated receptor-like kinase (SARK) show an increased tolerance to drought stress, attributable in part to CK-induced delayed leaf senescence (155). However, CK-deficient mutants in *Arabidopsis* also show strong stress tolerance to drought and salinity as well as ABA hypersensitivity and reduced levels of ABA (140). These results suggest finely tuned mechanisms for maintaining CK homeostasis and action, in part through adjustments in ABA synthesis and signaling.

CKs appear to contribute to the successful establishment of biotrophic infections (192). Recently, it was reported that the *Pst* effector HopQ1 suppresses immune responses in *Arabidopsis* and induces CK signaling in the plant (72). However, CKs may also interact with SA through their activation of ARR2 TFs that participate in SA-mediated defense gene expression (37) (**Figure 6c**). How the plant negotiates these apparent countervailing CK actions is unclear, although the levels of CKs may be the critical factor here, with low concentrations promoting susceptibility and high concentrations promoting resistance (72). Although the adaptive role of CKs in plants under drought and salinity stress is complicated, there is intense interest in how plants balance the stress-adaptive functions of ABA and CK in efforts to improve environmental stress tolerance, which may have implications for tolerance to pathogens as well (112, 199).

DELLA Proteins and Gibberellic Acid

DELLA proteins are a nuclear family of TFs that are now recognized as integrators of phytohormone signaling. DELLAs repress growth and GA signaling, and GA reciprocates by derepressing its pathway to promote degradation of DELLAs via the ubiquitin-proteasome system (190) (**Figure 6d**). In *Arabidopsis*, there are five DELLA genes that encode proteins directed at different and overlapping targets in processes associated with germination, cellular redox, growth, and abiotic and biotic stress responses, and mutations in these genes lead to a number of altered GA and stress phenotypes (67, 190). One of these DELLA proteins, RGL2, is important in repressing seed germination and is induced by ABA, and in turn increases ABA levels and the expression of the TF ABI5. In addition to their role as repressors of GA signaling and of growth, DELLAs are stabilized by ABA and ET to potentiate cross talk between SA and JA through promoting JA action and inhibiting SA synthesis and signaling (67, 137). JA's inhibition of plant growth in favor

of defense appears to be attributable in large part to interference with GA signaling that leads to DELLA protein degradation, indicating antagonism between JA and GA (204). DELLAs bind to JAZ repressor proteins to liberate MYC2 and promote JA-responsive gene expression. DELLAs mediate salt stress-induced inhibition of growth, and an *Arabidopsis* mutant with knockouts of four of the DELLA genes is more resistant to salt than are wild-type plants (1). In *Arabidopsis*, DELLAs also increase susceptibility to *Pst* DC3000 (137). Likewise, the quadruple DELLA knockout shows increased resistance to *Pst* DC3000, generally regarded as a hemibiotroph, but increased susceptibility to the necrotrophic fungi *Alternaria brassicicola* (137) and *Botrytis cinerea* (2). The evidence positions DELLAs as potentially important contributors to the predisposing effects of salt and other stresses on disease, and further research is needed to address this issue.

Auxin

Auxin integrates the effects of multiple phytohormones to regulate plant growth and development and root system architecture in response to changing environments (64, 82). Auxin stimulates lateral root formation as a response to nutrient limitation; however, drought and salinity stress repress auxin signaling and lateral root development to favor primary root growth to tap water deeper in the soil (70). The interplay of auxin and the counteracting effects of ABA and CK contribute to these different outcomes, due in part to differential regulation of the TF ABI4 and adjustments to polar auxin transport (170).

Auxin's role in plant-microbe interactions is well established in the traditional sense, with pathogen- and host-derived IAA contributing to disease phenotypes such as galls and fruit russetting. However, auxin action in predisposition has not been thoroughly investigated, although there is compelling evidence for auxin modulating SA and JA defense networks at several levels (**Figure 6e**). The major categories of auxin-responsive genes are grouped in the *Aux/IAA*, *GH3*, and small auxin-up RNAs (SAUR) families. *AUX/IAA* proteins are transcriptional repressors that are removed from their targets by auxin-activated SCF^{TIR1} receptor complexes and tagged for proteasome degradation by the SCF E3 ubiquitin ligase pathway (20). In *Arabidopsis*, the eubacterial PAMP flg22 binds to the receptor kinase FLS2 to trigger an immune response that includes posttranscriptional silencing of mRNAs for the F-box auxin receptors TIR1, AFB2, and AFB3 (138). This stabilizes *Aux/IAA* repressors, and the repression in auxin signaling further enhances resistance to *Pst* DC3000. SA treatment also downregulates auxin signaling in a similar fashion (194), providing evidence that auxin and SA counteract each other, with auxin diminishing resistance and SA having the opposite effect, at least on *Pst*. In contrast to *Pst*, necrotrophic fungal pathogens exploit auxin signaling during pathogenesis by stabilizing another group of auxin transcriptional repressors (i.e., AXRs), resulting in enhanced susceptibility (108). The different outcomes might be explained by *Arabidopsis*'s contrasting defense strategies in dealing with biotrophs and necrotrophs (66).

Another level of control of auxin action is provided by the auxin-inducible *GH3* genes, which encode acyl acid amido synthetases that regulate the endogenous auxin pool by catalyzing formation of auxin-amino acid conjugates (144). One of these, *WES1*, displays complex regulation and induction by SA, ABA, and *Pst* infection as well as drought, cold, and heat stresses. An over-expressing mutant (*wes1-D*) displays severe dwarfing characteristic of auxin deficiency as well as altered phenotypes to the various abiotic and biotic stresses. Other *GH3* family members encode enzymes that conjugate amino acids to JA (i.e., JA-Ile) and SA, providing another mechanism for adjusting levels of these phytohormones and cross talk (173).

An intriguing model positions auxin as a transitional signal between phases of JA and SA signaling to establish systemic acquired resistance (SAR) (186), predicated in part on the above-cited

studies but also on the key observation of an early transient spike in JA that precedes SA signaling in *Arabidopsis* following inoculation with incompatible *Pst*. In this model, the initial JA phase induces indole biosynthesis that includes auxin as well as JAZ repressors to begin to dampen JA signaling. As JA action subsides, the auxin phase follows and engages the auxin importer AUX1 to import and position IAA for interaction with the TIR receptor to derepress auxin-regulated transcription. Auxin induction of *GH3* genes provides a mechanism to diminish auxin signaling, facilitating the next transition to SA-mediated defense activation and priming for SAR. It will be interesting to see if this model can be generalized to other host-pathogen interactions and whether these signaling transitions are disrupted by ABA and predisposing abiotic stresses.

Brassinosteroids

BRs, e.g., brassinolide, are a group of more than 70 polyhydroxysteroids with impacts on various aspects of plant growth and development as well as a potential protective role in abiotic and biotic stress responses (142). Application of BRs can induce protection against osmotic, drought, and temperature stresses, suggesting that they could counter predisposition to diseases. BRs induce oxo-phytodienoic reductase in JA biosynthesis and when applied with other phytohormones also show additive or synergistic effects (e.g., GA and auxin) or mutual antagonism (e.g., ABA) (209). The impact of BR treatment on host-pathogen interactions is mixed, with examples of increased resistance (135) as well as increased susceptibility (16). BR signaling has emerged as a potential target for engineering tolerance to both abiotic and biotic stresses (52).

BR perception and signaling are tightly regulated through phosphorylation and dephosphorylation reactions (**Figure 6f**). The BR INSENSITIVE 1 (BRI1) receptor is a leucine-rich repeat receptor kinase (LRR-RK) that requires BAK1 (BRI1-associated kinase) as a coreceptor to positively regulate downstream BR signaling. An interesting connection between BR and PAMP-triggered immunity occurs at the level of their perception by RKs. The PAMP receptor FLS2 almost immediately after binding flg22 forms a heteromer with BAK1 to trigger transphosphorylation and a phosphorylation cascade of downstream targets (178). Although FLS2 and BRI1 would appear to negatively regulate the other's action by competing for BAK1, experimental evidence indicates that the outcome of their interaction can be positive or negative, depending on levels of BR, BRI1, and BAK1 (196). The evidence points to coordination of BR signaling and PAMP-triggered immunity during growth and abiotic/biotic stress events. Additional details of receptor-like kinases and BR perception and response are reviewed in Osakabe et al. (142).

Oxidative Stress and Reactive Oxygen Species Signaling in Predisposition

ROS generation is an initiating event in biotic and abiotic stress responses, and ROS signaling is integrated with multiple phytohormone networks (126). ROS function as both localized and distance signals through their propagation as waves in the plant from an initiating stimulus. Phosphorelay systems are important both in generating ROS and in modulating ROS signaling. ROS generation by various isoforms of the *Arabidopsis* NADPH-dependent respiratory burst oxidase homolog oxidase complex (i.e., AtRBOH) involves calcium signaling and phosphorylation (126). In turn, MAP kinase cascades are engaged by elevated ROS levels to regulate TFs and gene expression, leading to stress adaptation, such as cold- and salt-stress tolerance (182). A MAPK cascade that couples ABA, catalase induction, and H₂O₂ generation is mediated by MKK1 and MPK6 (200).

Tight regulation of the steady-state levels of ROS is required in multiple cellular processes in plants (8), with more than 150 genes in the network managing ROS in *Arabidopsis* (125). Analysis of

the NaCl-induced transcriptome in roots suggests a high degree of subfunctionalization within this network at the regulatory and catalytic levels (85). The majority of genes in the ROS-scavenging network are unresponsive or downregulated within the salt-stress regimes we have found to be capable of predisposing. Water stress induces changes in ROS levels, and many ABA-regulated genes are also induced by oxidative stress (36). *AtrbobD* and *AtrbobF* appear to be required for ABA-induced ROS generation and are linked to ABA-induced changes in guard cell turgor and stomatal function as well as hypersensitive cell death in response to avirulent pathogens (160). ABA and water stress upregulate transcripts of abscisic aldehyde oxidase (AO; e.g., *AAO3*) (Figure 7) and transcripts for *LOS5/ABA3*, responsible for AO and xanthine dehydrogenase (XDH) activation (207). Thus, water stress can enhance ROS production via AO and XDH in an ABA-dependent manner, and place Rboh, AO, and XDH as candidates for ABA-dependent ROS transducers involved in predisposition.

Transcription Factors and Network Interactions

There are a number of important TFs not mentioned above that provide points of convergence in abiotic and biotic stress response networks and have the potential to alter disease outcomes during predisposing stress (63, 203). Atkinson & Urwin (13) discussed TF functions more from the point of view of coordinating gene expression to enhance stress tolerance and defense against pathogens. Notable among these are various members belonging to the MYB, NAC, AREB/ABF (ABA response-element binding factor), GBFs (G-box binding factors), and AP2/ERF families that are regulated by ABA. MYB family members are important in gene expression for the biosynthesis of phenylpropanoids, flavonols, and cuticular wax as well as other defensive products, including ABA-dependent SA accumulation, and seem to enable the plant to discriminate between different stress signals (164). NAC family members, such as ATAF1, ATAF2, and RD26 in *Arabidopsis*, are inducible by ABA and various abiotic and biotic stresses (84). The AREB/ABFs and GBFs are subgroups within the bZIP class of TFs and are important integrators of ABA signal-response coupling (77). AP2/ERF family members were discussed in relation to ET signaling, but others, such as DEAR1 (*Arabidopsis*) and TS1 (tobacco), contribute to abiotic and biotic stress regulation of gene expression. Overexpression of *TS1* enhances tolerance to osmotic stress and resistance to *Pseudomonas syringae* pv. *tabaci* (145). DEAR1 represses *DREB* genes to diminish tolerance to cold stress but enhances pathogen resistance (187). The *DREB/CBF* TFs are ERF family members that bind to *cis*-acting promoter elements of genes conferring dehydration and cold tolerance (5). In *Arabidopsis*, *DREB1A* is induced by cold but not by drought or ABA. ABA-independent targets should also be considered in predisposition studies.

PROGRAMMED CELL DEATH AND PREDISPOSITION

The connection of PCD with abiotic stress and disease is indicated by hallmark PCD features in cells responding to various stresses and the broad-spectrum protection afforded by transgenic expression of antiapoptotic genes (49). Expression of proapoptotic (e.g., *Bax1*) and antiapoptotic genes (e.g., *Bcl-2*, *CED9*, *P35*, *AtB11*) from diverse organisms respectively induce and protect plants against PCD, providing evidence for cross-kingdom commonalities in cell death mechanisms. For example, SflAP, a negative regulator of PCD from the insect *Spodoptera frugiperda*, is a member of the IAP (inhibitor of apoptosis) family of proteins (49). Transgenic expression of SflAP in tobacco and tomato plants confers tolerance to heat, salt, and the mycotoxin FB1 as well as resistance to the necrotrophic fungal pathogens *A. alternata* and *Sclerotinia sclerotiorum* (107). Also, SflAP-transgenic tomatoes are impaired in *EIN3* and ET-mediated gene expression, display delayed fruit

ripening, and are less sensitive to the inhibitory effect of ABA on plant growth. SflAP appears to have a conserved function for inhibiting stress-induced cell death, and when expressed in plants may operate through ABA- and ET-regulated mechanisms. SflAP's protection is attributable to its E3 ubiquitin ligase activity, and regulated protein degradation via the proteasome is essential for SflAP-induced stress tolerance and delayed fruit ripening (87). Targeting PCD is an attractive strategy with the potential for improving abiotic and biotic stress tolerance in plants. Ensuring tight regulation to avoid unwanted effects on plant growth and development and understanding the limitations of PCD within different biological and environmental contexts will be crucial for deployment.

SMALL RNAs IN ABIOTIC STRESS AND DISEASE

Small RNAs (sRNA) play various roles in plant processes, such as developmental patterning and genome integrity, but are best known for RNA silencing in plant-virus interactions (14). These short, 18–25 nucleotide sequences are diverse in genomic distribution, sequence, biogenesis, and regulatory function. MicroRNAs (miRNAs) are a class of sRNAs that target complementary mRNAs for translational repression or cleave their targets through association with RNA-induced silencing complexes (RISCs). Emerging evidence indicates important roles for miRNAs in abiotic stress responses (179). The first implication of sRNAs in ABA signaling was from experiments with the *Arabidopsis hyl1* mutant, which is hypersensitive to ABA. HYL1 partners with DICER LIKE 1 (DCL1) for the biogenesis and accumulation of miRNAs but is not required for posttranscriptional gene regulation (110). The first ABA-induced miRNA to be discovered is *miR393*, which is also strongly upregulated by dehydration, increased salinity, and cold (179). Subsequently, *miR393* was found to be induced by *flg22* and to repress auxin signaling as described above (138). *miR159* is also induced by ABA and targets two MYB TFs that are positive regulators of ABA responses in *Arabidopsis* (147).

Plants may rely on miRNA-mediated degradation of ABA signaling factors following stress episodes in order to reduce ABA levels and restore the prestress physiological state (154). In *Arabidopsis*, miRNAs upregulated by ABA include *miR160*, *miR417*, and *miR319*; ABA-downregulated miRNAs include *miR167*, *miR169*, and *miR398* (91). The sets of miRNAs associated with salt and drought stress both overlap with and differ from those induced by ABA, with some showing inverse responses to ABA and stress treatments (91). Similarities in the miRNA expression profiles between drought and salt stress and ABA strengthen a role for miRNAs in ABA-mediated stress responses. These miRNAs mainly target diverse TF families, such as SBPs (squamosa promoter binding protein), MYBs/TCPs (myeloblastosis/teosinte branched1, cycloidea, and PCF), ARFs (auxin response factor), HD-ZIPs (homeodomain leucine zipper), and NFY (nuclear transcription factor Y) subunits.

miRNA regulation in biotic stress responses is highly variable and is likely influenced by the host, the pathogen, and the tissue infected (91). Differential miRNA regulation has been reported in symbiotic interactions and in pathogenic viral, bacterial, fungal, and nematodal interactions. In the soybean–*Phytophthora sojae* interaction, infection altered the expression of a small group of host miRNAs with known roles in abiotic stresses (68). Many of the aforementioned stress-related miRNAs are also upregulated by bacterial pathogens and downregulated by fungal pathogens (91). Recent work has shown that many solanaceous species possess miRNAs of the *miR482/miR2118* superfamily. Members of this miRNA superfamily specifically target the P-loop motif of mRNAs that encode NBS-LRR resistance proteins (169). Degradation of the target mRNA results in the creation of phased, secondary small RNAs (phasRNAs), which allows basal expression of *miR482* to simultaneously silence multiple *R* genes (151). *miR482* is also suppressed by viral, bacterial,

and fungal pathogens, allowing R proteins to accumulate (169, 211), and is upregulated by abiotic stress (9). Deciphering complex small RNA regulatory networks will require the identification of sets of miRNAs and their targets. Such information will provide additional insights into how plants cope with concurrent biotic and abiotic stresses.

SUMMARY

During evolution, plants developed finely tuned mechanisms to cope with diverse abiotic and biotic stresses. This is our operational paradigm and there are many studies that affirm cooperative signaling and response to mitigate environmental challenges and enhance stress tolerance. Yet we must reconcile the fact that plants often succumb to pathogens that they would normally resist when confronted with transient and relatively mild abiotic stresses. Modern agricultural crops must balance disease and pest resistance, stress tolerance, and growth with associated fitness costs (43), but how well they do this can depend on selection criteria during breeding as well as on the impact of production practices designed to maximize yields or to conserve increasingly limited resources. The details of stress network interactions within current cultivars, which are often adapted to rich resource environments, may differ from land races and old world varieties and depart from the experimental models. The circuitry, although complex and interconnected, may not be as elegantly wired as we would wish to offset the unproductive interactions.

ABA has emerged as a global regulator of abiotic stress responses and an important phytohormone in plant-microbe interactions with systemic effects on resistance and susceptibility (Figures 6, 7, and 8). It has become apparent that cross talk occurs among all the major phytohormones during stress events, and the challenge is discerning which interactions most influence disease development in particular contexts. Transcriptomics has enabled identification of large sets of coordinately regulated stress genes, but interpretation of the results in the context of predisposition can be confounded in that the output may lack sufficient resolution to ascribe functional correspondence to disease outcomes. For example, many ROS- or ABA-regulated defense genes are induced in both susceptible and resistant interactions as well as by other stresses, but issues of timing and intensity as well as whether mRNAs are translated may not be apparent from the data without further analysis. Critical for interpretation of such studies are corroborative quantitative assays that rigorously distinguish disease phenotypes.

Delineation of stress network interactions has implications for disease management, such as in deploying chemically or biologically induced resistance (156, 193). The efficacy of plant activators depends on the environmental context, and abiotic stresses are expected to influence JA and SA network dynamics during pathogen attack (24). Encouraging are results acquired when using the SA mimic tiadinil, albeit conducted under highly controlled experimental conditions, which induces disease protection in tomato under salinity stress (152). This and other studies suggest that plant activators may offset or at least function adequately under conditions of predisposing stress, but further assessment under the environmental rigors of the field is needed.

Various strategies under consideration to enhance plant stress tolerance include using agonists and antagonists of phytohormone action, engineering phytohormone response pathways, and breeding (143, 146, 199). An interesting example of the latter is in soybean, where a quantitative trait loci analysis of recombinant inbred lines found that genes for both flooding tolerance and resistance to *P. sojae* were necessary to reduce disease under flooded conditions (139), illustrating the potential synergy derived from combining traits for stress tolerance and disease resistance. Regardless of the strategy, it is important to critically assess how plants modified for abiotic stress




Figure 8

General model and recent areas for mechanistic inquiry into the role of stress-induced abscisic acid (ABA) in predisposition. Abbreviations: BR, brassinosteroid; CK, cytokinin; ET, ethylene; GA, gibberellic acid; IAA, indole-acetic acid; JA, jasmonic acid; ROS, reactive oxygen species; SA, salicylic acid.

tolerance perform when challenged by pathogens and pests. An understanding of the adverse interactions among phytohormone networks in the face of concurrent abiotic and biotic stresses is emerging, and the challenge is to determine how and to what extent the underlying processes can be manipulated to effect positive outcomes.

SUMMARY POINTS

1. Mild, episodic stresses can severely predispose plants to pathogens they would otherwise resist.
2. Molecular and biochemical studies indicate extensive overlap in abiotic and biotic stress responses, with evidence for a universal stress response transcriptome (113). Understanding the degree to which there is fine-tuning to tailor responses to different stresses is an ongoing quest.
3. Strong evidence supports ABA's determinative role in predisposition, but there is also extensive cross talk among all the phytohormones in stress events.

4. A model for the recruitment of ROS and phytohormones to sequentially engage defense responses is emerging (186), but how the sequence is disrupted by predisposing stress events is unclear.
5. Strategies for improving stress tolerance in plants should also consider impacts on disease resistance.

FUTURE ISSUES

1. Comparative studies are needed to assess how stress network dynamics discovered in *Arabidopsis* translate to agricultural pathosystems.
2. Studies that compound multiple stresses are needed to fully characterize the abiotic-biotic stress interactome as it may occur in the field (124, 150) and likely will require a robust systems biology approach involving transcriptomics, proteomics, and metabolomics to identify key regulatory hubs and outputs (40, 158).
3. There is a need for high-throughput methods for phenotyping root-pathogen interactions.
4. There are more than 600 receptor-like kinases in *Arabidopsis* (142). Many appear to be involved in abiotic and biotic stress perception but require functional characterization.
5. Small RNAs, chromatin remodeling, and epigenetics are emerging areas in plant stress biology. How these processes impact disease outcomes is an active area of inquiry.

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

The authors are grateful to Professors Thomas R. Gordon and Martin B. Dickman for their critical reviews of the manuscript and to Dr. Matthew V. DiLeo and Professors James D. MacDonald, John M. Duniway, and David M. Rizzo for productive past collaborations. Appreciation is also extended to Professor Lincoln Taiz for permission to use graphics within **Figure 5**. We also acknowledge the many scientists whose papers we were unable to cite directly because of the space limits required by the publisher. Support for the authors' research on predisposition has been provided by various grants from the USDA NIFA (National Institute of Food and Agriculture), USDA Forest Service, USDA APHIS (Animal and Plant Health Inspection Service), and the University of California Agricultural Experiment Station.

LITERATURE CITED

1. Achard P, Cheng H, De Grauwe L, Decat J, Schoutteten H, et al. 2006. Integration of plant responses to environmentally activated phytohormonal signals. *Science* 311:91–94
2. Achard P, Renou JP, Berthome R, Harberd NP, Genschik P. 2008. Plant DELLAs restrain growth and promote survival of adversity by reducing the levels of reactive oxygen species. *Curr. Biol.* 18:656–60

3. Achuo EA, Prinsen E, Hofte M. 2006. Influence of drought, salt stress and abscisic acid on the resistance of tomato to *Botrytis cinerea* and *Oidium neolycopersici*. *Plant Pathol.* 55:178–86
4. Adie BAT, Perez-Perez J, Perez-Perez MM, Godoy M, Sanchez-Serrano JJ, et al. 2007. ABA is an essential signal for plant resistance to pathogens affecting JA biosynthesis and the activation of defenses in *Arabidopsis*. *Plant Cell* 19:1665–81
5. Agarwal PK, Agarwal P, Reddy MK, Sopory SK. 2006. Role of DREB transcription factors in abiotic and biotic stress tolerance in plants. *Plant Cell Rep.* 25:1263–74
6. Albacete A, Ghanem ME, Martinez-Andujar C, Acosta M, Sanchez-Bravo J, et al. 2008. Hormonal changes in relation to biomass partitioning and shoot growth impairment in salinized tomato (*Solanum lycopersicum* L.) plants. *J. Exp. Bot.* 59:4119–31
7. Anderson JP, Badruzaufari E, Schenk PM, Manners JM, Desmond OJ, et al. 2004. Antagonistic interaction between abscisic acid and jasmonate-ethylene signaling pathways modulates defense gene expression and disease resistance in *Arabidopsis*. *Plant Cell* 16:3460–79
8. Apel K, Hirt H. 2004. Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annu. Rev. Plant Biol.* 55:373–99
9. Arenas-Huertero C, Pérez B, Rabanal F, Blanco-Melo D, De la Rosa C, et al. 2009. Conserved and novel miRNAs in the legume *Phaseolus vulgaris* in response to stress. *Plant Mol. Biol.* 70:385–401
10. Asselbergh B, Achuo AE, Hofte M, Van Gijsegem F. 2008. Abscisic acid deficiency leads to rapid activation of tomato defence responses upon infection with *Erwinia chrysanthemi*. *Mol. Plant Pathol.* 9:11–24
11. Asselbergh B, Curvers K, Franca SC, Audenaert K, Vuylsteke M, et al. 2007. Resistance to *Botrytis cinerea* in *sitiens*, an abscisic acid-deficient tomato mutant, involves timely production of hydrogen peroxide and cell wall modifications in the epidermis. *Plant Physiol.* 144:1863–77
12. Asselbergh B, De Vleeschauwer D, Hofte M. 2008. Global switches and fine-tuning-ABA modulates plant pathogen defense. *Mol. Plant-Microbe Interact.* 21:709–19
13. Atkinson NJ, Urwin PE. 2012. The interaction of plant biotic and abiotic stresses: from genes to the field. *J. Exp. Bot.* 63:3523–43
14. Axtell MJ. 2013. Classification and comparison of small RNAs from plants. *Annu. Rev. Plant Biol.* 64:137–59
15. Ayres PG. 1984. The interaction between environmental stress injury and biotic disease physiology. *Annu. Rev. Phytopathol.* 22:53–75
16. Bajguz A, Hayat S. 2009. Effects of brassinosteroids on the plant responses to environmental stresses. *Plant Physiol. Biochem.* 47:1–8
17. Barkosky RR, Einhellig FA. 1993. Effects of salicylic acid on plant water relationships. *J. Chem. Ecol.* 19:237–47
18. Belkhadir Y, Jaillais Y, Epple P, Balsemao-Pires E, Dangl JL, Chory J. 2012. Brassinosteroids modulate the efficiency of plant immune responses to microbe-associated molecular patterns. *Proc. Natl. Acad. Sci. USA* 109:297–302
19. Bender J, Weigel HJ. 2011. Changes in atmospheric chemistry and crop health: a review. *Agron. Sustain. Dev.* 31:81–89
20. Benjamins R, Scheres B. 2008. Auxin: the looping star in plant development. *Annu. Rev. Plant Biol.* 59:443–65
21. Bettucci L, Alonso R, Tiscornia S. 1999. Endophytic mycobiota of healthy twigs and the assemblage of species associated with twig lesions of *Eucalyptus globulus* and *E. grandis* in Uruguay. *Mycol. Res.* 103:468–72
22. Birkenbihl RP, Diezel C, Somssich IE. 2012. *Arabidopsis* WRKY33 is a key transcriptional regulator of hormonal and metabolic responses toward *Botrytis cinerea* infection. *Plant Physiol.* 159:266–85
23. Blaker NS, Macdonald JD. 1986. The role of salinity in the development of *Phytophthora* root rot of citrus. *Phytopathology* 76:970–75
24. Bostock RM. 2005. Signal crosstalk and induced resistance: straddling the line between cost and benefit. *Annu. Rev. Phytopathol.* 43:545–80
25. Bostock RM, Quatrano RS. 1992. Regulation of *Em* gene expression in rice: interaction between osmotic stress and abscisic acid. *Plant Physiol.* 98:1356–63

26. Boter M, Ruiz-Rivero O, Abdeen A, Prat S. 2004. Conserved MYC transcription factors play a key role in jasmonate signaling both in tomato and *Arabidopsis*. *Genes Dev.* 18:1577–91
27. Boyer JS. 1995. Biochemical and biophysical aspects of water deficits and the predisposition to disease. *Annu. Rev. Phytopathol.* 33:251–74
28. Brossa R, Lopez-Carbonell M, Jubany-Mari T, Alegre L. 2011. Interplay between abscisic acid and jasmonic acid and its role in water-oxidative stress in wild-type, ABA-deficient, JA-deficient, and ascorbate-deficient *Arabidopsis* plants. *J. Plant Growth Regul.* 30:322–33
29. Bultreys A, Kaluzna M. 2010. Bacterial cankers caused by *Pseudomonas syringae* on stone fruit species with special emphasis on the pathovars *syringae* and *morsprunorum* race 1 and race 2. *J. Plant Pathol.* 92:S21–33
30. Burgess T, McComb JA, Colquhoun I, Hardy GES. 1999. Increased susceptibility of *Eucalyptus marginata* to stem infection by *Phytophthora cinnamomi* resulting from root hypoxia. *Plant Pathol.* 48:797–806
31. Cahill DM, Ward EWB. 1989. Rapid localized changes in abscisic acid concentrations in soybean in interactions with *Phytophthora megasperma* f. sp. *glycinea* or after treatment with elicitors. *Physiol. Mol. Plant Pathol.* 35:483–94
32. Cao FY, Yoshioka K, Desveaux D. 2011. The roles of ABA in plant-pathogen interactions. *J. Plant Res.* 124:489–99
33. Chakraborty S. 2011. Climate change and plant diseases. *Plant Pathol.* 60:2–14
34. Chan ZL. 2012. Expression profiling of ABA pathway transcripts indicates crosstalk between abiotic and biotic stress responses in *Arabidopsis*. *Genomics* 100:110–15
35. Chapin FS. 1991. Integrated responses of plants to stress. *Bioscience* 41:29–36
36. Cho D, Shin DJ, Jeon BW, Kwak JM. 2009. ROS-mediated ABA signaling. *J. Plant Biol.* 52:102–13
37. Choi J, Choi D, Lee S, Ryu CM, Hwang I. 2011. Cytokinins and plant immunity: old foes or new friends? *Trends Plant Sci.* 16:388–94
38. Christmann A, Grill E, Huang J. 2013. Hydraulic signals in long-distance signaling. *Curr. Opin. Plant Biol.* 16:293–300
39. Cook RJ, Papendick RI. 1972. Influence of water potential of soils and plants on root disease. *Annu. Rev. Phytopathol.* 10:349–74
40. Cramer GR, Urano K, Delrot S, Pezzotti M, Shinozaki K. 2011. Effects of abiotic stress on plants: a systems biology perspective. *BMC Plant Biol.* 11:163
41. Cutler SR, Rodriguez PL, Finkelstein RR, Abrams SR. 2010. Abscisic acid: emergence of a core signaling network. *Annu. Rev. Plant Biol.* 61:651–79
42. Davison EM, Tay FCS. 1987. The effect of waterlogging on infection of *Eucalyptus marginata* seedlings by *Phytophthora cinnamomi*. *New Phytol.* 105:585–94
43. Denance N, Sanchez-Vallet A, Goffner D, Molina A. 2013. Disease resistance or growth: the role of plant hormones in balancing immune responses and fitness costs. *Front. Plant Sci.* 4:155
44. de Ollas C, Hernando B, Arbona V, Gomez-Cadenas A. 2013. Jasmonic acid transient accumulation is needed for abscisic acid increase in citrus roots under drought stress conditions. *Physiol. Plant* 147:296–306
45. Desprez-Loustau ML, Marcais B, Nageleisen LM, Piuu D, Vannini A. 2006. Interactive effects of drought and pathogens in forest trees. *Ann. For. Sci.* 63:597–612
46. de Torres Zabala M, Bennett MH, Truman WH, Grant MR. 2009. Antagonism between salicylic and abscisic acid reflects early host-pathogen conflict and moulds plant defence responses. *Plant J.* 59:375–86
47. de Torres-Zabala M, Truman W, Bennett MH, Lafforgue G, Mansfield JW, et al. 2007. *Pseudomonas syringae* pv. *tomato* hijacks the *Arabidopsis* abscisic acid signalling pathway to cause disease. *EMBO J.* 26:1434–43
48. Dhindsa RS, Cleland RE. 1975. Water stress and protein synthesis. I. Differential inhibition of protein synthesis. *Plant Physiol.* 55:778–81
49. Dickman MB, Fluhr R. 2013. Centrality of host cell death in plant-microbe interactions. *Annu. Rev. Phytopathol.* 51:543–70
50. DiLeo MV, Pye MF, Roubtsova TV, Duniway JM, MacDonald JD, et al. 2010. Abscisic acid in salt stress predisposition to *Phytophthora* root and crown rot in tomato and *Chrysanthemum*. *Phytopathology* 100:871–79

51. Dinneny JR, Long TA, Wang JY, Jung JW, Mace D, et al. 2008. Cell identity mediates the response of *Arabidopsis* roots to abiotic stress. *Science* 320:942–45
52. Divi UK, Rahman T, Krishna P. 2010. Brassinosteroid-mediated stress tolerance in *Arabidopsis* shows interactions with abscisic acid, ethylene and salicylic acid pathways. *BMC Plant Biol.* 10:151
53. Dordas C. 2008. Role of nutrients in controlling plant diseases in sustainable agriculture. A review. *Agron. Sustain. Dev.* 28:33–46
54. Dreher K, Callis J. 2007. Ubiquitin, hormones and biotic stress in plants. *Ann. Bot.* 99:787–822
55. Duke SO, Wedge DE, Cerdeira AL, Matallo MB. 2007. Interaction of synthetic herbicides with plant disease and microbial herbicides. In *Novel Biotechnologies for Biocontrol Agent Enhancement and Management*, ed. M Vurro, J Gressel, pp. 277–96. Dordrecht, Neth.: Springer
56. Duniway JM. 1977. Predisposing effect of water stress on severity of *Phytophthora* root rot in safflower. *Phytopathology* 67:884–89
57. Duniway JM. 1979. Water relations of water molds. *Annu. Rev. Phytopathol.* 17:431–60
58. Erbaugh DK, Windham MT, Stodola AJW, Auge RM. 1995. Light intensity and drought stress as predisposition factors for dogwood anthracnose. *J. Environ. Hort.* 13:186–89
59. Eyles A, Bonello P, Ganley R, Mohammed C. 2010. Induced resistance to pests and pathogens in trees. *New Phytol.* 185:893–908
60. Fan J, Hill L, Crooks C, Doerner P, Lamb C. 2009. Abscisic acid has a key role in modulating diverse plant-pathogen interactions. *Plant Physiol.* 150:1750–61
61. Finkelstein R. 2013. Abscisic acid synthesis and response. *Arabidopsis Book* 11:e0166
62. Fujita M, Fujita Y, Noutoshi Y, Takahashi F, Narusaka Y, et al. 2006. Crosstalk between abiotic and biotic stress responses: a current view from the points of convergence in the stress signaling networks. *Curr. Opin. Plant Biol.* 9:436–42
63. Fujita Y, Fujita M, Shinozaki K, Yamaguchi-Shinozaki K. 2011. ABA-mediated transcriptional regulation in response to osmotic stress in plants. *J. Plant Res.* 124:509–25
64. Galvan-Ampudia CS, Testerink C. 2011. Salt stress signals shape the plant root. *Curr. Opin. Plant Biol.* 14:296–302
65. Garrett KA, Dendy SP, Frank EE, Rouse MN, Travers SE. 2006. Climate change effects on plant disease: genomes to ecosystems. *Annu. Rev. Phytopathol.* 44:489–509
66. Glazebrook J. 2005. Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens. *Annu. Rev. Phytopathol.* 43:205–27
67. Grant MR, Jones JDG. 2009. Hormone (dis)harmony moulds plant health and disease. *Science* 324:750–52
68. Guo N, Ye W-W, Wu X-L, Shen D-Y, Wang Y-C, et al. 2011. Microarray profiling reveals microRNAs involving soybean resistance to *Phytophthora sojae*. *Genome* 54:954–58
69. Guo W-L, Chen R-G, Gong Z-H, Yin Y-X, Li D-W. 2013. Suppression subtractive hybridization analysis of genes regulated by application of exogenous abscisic acid in pepper plant (*Capsicum annuum* L.) leaves under chilling stress. *PLoS ONE* 8:e66667
70. Ha S, Vankova R, Yamaguchi-Shinozaki K, Shinozaki K, Tran LSP. 2012. Cytokinins: metabolism and function in plant adaptation to environmental stresses. *Trends Plant Sci.* 17:172–79
71. Hahlbrock K, Chappell J, Jahnen W, Walter M. 1985. Early defense reactions of plants to pathogens. In *Molecular Form and Function of the Plant Genome*, ed. L van Vloten-Doting, GSP Groot, TC Hall, pp. 129–40. New York: Plenum Publ. Corp.
72. Hann DR, Dominguez-Ferreras A, Motyka V, Dobrev PI, Schornack S, et al. 2014. The *Pseudomonas* type III effector HopQ1 activates cytokinin signaling and interferes with plant innate immunity. *New Phytol.* 201:585–98
73. Hartig R. 1894. *Text-Book of the Diseases of Trees*. London: MacMillan and Co.
74. Hauser F, Waadt R, Schroeder JI. 2011. Evolution of abscisic acid synthesis and signaling mechanisms. *Curr. Biol.* 21:R346–55
75. Henfling J, Bostock R, Kuć J. 1980. Effect of abscisic acid on rishitin and lubimin accumulation and resistance to *Phytophthora infestans* and *Cladosporium cucumerinum* in potato tuber tissue slices. *Phytopathology* 70:1074–78

76. Heritage AD, Harrigan EKS. 1984. Environmental factors influencing safflower screening for resistance to *Phytophthora cryptogea*. *Plant Dis.* 68:767–69
77. Himmelbach A, Yang Y, Grill E. 2003. Relay and control of abscisic acid signaling. *Curr. Opin. Plant Biol.* 6:470–79
78. Horvath E, Szalai G, Janda T. 2007. Induction of abiotic stress tolerance by salicylic acid signaling. *J. Plant Growth Regul.* 26:290–300
79. Huang X, Li J, Bao F, Zhang X, Yang S. 2010. A gain-of-function mutation in the *Arabidopsis* disease resistance gene *RPP4* confers sensitivity to low temperature. *Plant Physiol.* 154:796–809
80. Huber DM, Graham RD. 1999. The role of nutrition in crop resistance and tolerance to disease. In *Mineral Nutrition of Crops: Fundamental Mechanisms and Implications*, ed. Z Rengel, pp. 205–26. New York: Food Prod. Press
81. Jackson MB. 2002. Long-distance signalling from roots to shoots assessed: the flooding story. *J. Exp. Bot.* 53:175–81
82. Jaillais Y, Chory J. 2010. Unraveling the paradoxes of plant hormone signaling integration. *Nat. Struct. Mol. Biol.* 17:642–45
83. Jakab G, Ton J, Flors V, Zimmerli L, Metraux JP, Mauch-Mani B. 2005. Enhancing *Arabidopsis* salt and drought stress tolerance by chemical priming for its abscisic acid responses. *Plant Physiol.* 139:267–74
84. Jensen MK, Lindemose S, de Masi F, Reimer JJ, Nielsen M, et al. 2013. ATAF1 transcription factor directly regulates abscisic acid biosynthetic gene *NCED3* in *Arabidopsis thaliana*. *FEBS Open Bio* 3:321–27
85. Jiang YQ, Deyholos MK. 2006. Comprehensive transcriptional profiling of NaCl-stressed *Arabidopsis* roots reveals novel classes of responsive genes. *BMC Plant Biol.* 6:25
86. Joshi-Saha A, Valon C, Leung J. 2011. A brand new START: abscisic acid perception and transduction in the guard cell. *Sci. Signal.* 4(201):re4
87. Kabbage M, Li W, Chen SR, Dickman MB. 2010. The E3 ubiquitin ligase activity of an insect anti-apoptotic gene (*SfIAP*) is required for plant stress tolerance. *Physiol. Mol. Plant Pathol.* 74:351–62
88. Kazan K, Manners JM. 2012. JAZ repressors and the orchestration of phytohormone crosstalk. *Trends Plant Sci.* 17:22–31
89. Kazan K, Manners JM. 2013. MYC2: the master in action. *Mol. Plant* 6:686–703
90. Keim R, Webster RK. 1974. Nitrogen fertilization and severity of stem rot of rice. *Phytopathology* 64:178–83
91. Khraiweh B, Zhu J-K, Zhu J. 2012. Role of miRNAs and siRNAs in biotic and abiotic stress responses of plants. *Biochim. Biophys. Acta* 1819:137–48
92. Kilian J, Peschke F, Berendzen KW, Harter K, Wanke D. 2012. Prerequisites, performance and profits of transcriptional profiling the abiotic stress response. *Biochim. Biophys. Acta* 1819:166–75
93. Kim Y, Park S, Gilmour SJ, Thomashow MF. 2013. Roles of CAMTA transcription factors and salicylic acid in configuring the low-temperature transcriptome and freezing tolerance of *Arabidopsis*. *Plant J.* 75:364–76
94. Koga H, Dohi K, Mori M. 2004. Abscisic acid and low temperatures suppress the whole plant-specific resistance reaction of rice plants to the infection of *Magnaporthe grisea*. *Physiol. Mol. Plant Pathol.* 65:3–9
95. Kohli A, Sreenivasulu N, Lakshmanan P, Kumar PP. 2013. The phytohormone crosstalk paradigm takes center stage in understanding how plants respond to abiotic stresses. *Plant Cell Rep.* 32:945–57
96. Kozłowski TT, Pallardy SG. 2002. Acclimation and adaptive responses of woody plants to environmental stresses. *Bot. Rev.* 68:270–334
97. Krasensky J, Jonak C. 2012. Drought, salt, and temperature stress-induced metabolic rearrangements and regulatory networks. *J. Exp. Bot.* 63:1593–608
98. Kriesel K. 1997. Effect of growth regulators on the susceptibility of pine seedlings to *Cylindrocarpum destructans* (Zins.) Scholt. and on the pathogenicity of this fungus to these seedlings: Part II. Abscisic acid (ABA). *Acta Univ. Nicolai Copernici Biol.* 52:45–50
99. Kuan TL, Erwin DC. 1980. Predisposition effect of water saturation of soil on *Phytophthora* root rot of alfalfa. *Phytopathology* 70:981–86
100. Kumar M, Busch W, Birke H, Kemmerling B, Nuernberger T, Schoeffl F. 2009. Heat shock factors HsfB1 and HsfB2b are involved in the regulation of *Pdfl.2* expression and pathogen resistance in *Arabidopsis*. *Mol. Plant* 2:152–65

101. Kuwabara C, Imai R. 2009. Molecular basis of disease resistance acquired through cold acclimation in overwintering plants. *J. Plant Biol.* 52:19–26
102. Leon-Reyes A, Du YJ, Koornneef A, Proietti S, Korbes AP, et al. 2010. Ethylene signaling renders the jasmonate response of *Arabidopsis* insensitive to future suppression by salicylic acid. *Mol. Plant-Microbe Interact.* 23:187–97
103. Leon-Reyes A, Spoel SH, De Lange ES, Abe H, Kobayashi M, et al. 2009. Ethylene modulates the role of NONEXPRESSOR OF PATHOGENESIS-RELATED GENES1 in cross talk between salicylate and jasmonate signaling. *Plant Physiol.* 149:1797–809
104. Leshem YY, Kuiper PJ, Erdei L, Lurie S, Perl-Treves R. 1998. Do Selye’s mammalian “GAS” concept and “co-stress” response exist in plants? *Ann. N.Y. Acad. Sci.* 851:199–208
105. Li BH, Li Q, Xiong LM, Kronzucker HJ, Kramer U, Shi WM. 2012. *Arabidopsis* plastid AMOS1/EGY1 integrates abscisic acid signaling to regulate global gene expression response to ammonium stress. *Plant Physiol.* 160:2040–51
106. Li J, Brader G, Kariola T, Palva ET. 2006. WRKY70 modulates the selection of signaling pathways in plant defense. *Plant J.* 46:477–91
107. Li W, Kabbage M, Dickman MB. 2010. Transgenic expression of an insect inhibitor of apoptosis gene, *SfLAP*, confers abiotic and biotic stress tolerance and delays tomato fruit ripening. *Physiol. Mol. Plant Pathol.* 74:363–75
108. Llorente F, Muskett P, Sanchez-Vallet A, Lopez G, Ramos B, et al. 2008. Repression of the auxin response pathway increases *Arabidopsis* susceptibility to necrotrophic fungi. *Mol. Plant* 1:496–509
109. Lopez-Zamora I, Bliss C, Jokela EJ, Comerford NB, Grunwald S, et al. 2007. Spatial relationships between nitrogen status and pitch canker disease in slash pine planted adjacent to a poultry operation. *Environ. Pollut.* 147:101–11
110. Lu C, Fedoroff N. 2000. A mutation in the *Arabidopsis* *HYL1* gene encoding a dsRNA binding protein affects responses to abscisic acid, auxin, and cytokinin. *Plant Cell* 12:2351–65
111. Lyzenga WJ, Stone SL. 2012. Abiotic stress tolerance mediated by protein ubiquitination. *J. Exp. Bot.* 63:599–616
112. Ma QH. 2008. Genetic engineering of cytokinins and their application to agriculture. *Crit. Rev. Biotechnol.* 28:213–32
113. Ma SS, Bohnert HJ. 2007. Integration of *Arabidopsis thaliana* stress-related transcript profiles, promoter structures, and cell-specific expression. *Genome Biol.* 8:R49
114. Ma SS, Gong QQ, Bohnert HJ. 2006. Dissecting salt stress pathways. *J. Exp. Bot.* 57:1097–107
115. MacDonald JD. 1982. Effect of salinity stress on the development of *Phytophthora* root rot of chrysanthemum *Chrysanthemum morifolium* cultivar Paragon. *Phytopathology* 72:214–19
116. MacDonald JD. 1984. Salinity effects on the susceptibility of chrysanthemum roots to *Phytophthora cryptogea*. *Phytopathology* 74:621–24
117. MacDonald JD. 1991. Heat stress enhances *Phytophthora* root rot severity in container-grown chrysanthemums. *J. Am. Soc. Hortic. Sci.* 116:36–41
118. Marek SM, Yaghmour M, Bostock RM. 2013. *Fusarium* spp., *Cylindrocarpon* spp., and predisposing environmental stress in the etiology of a canker disease of cold-stored fruit and nut tree seedlings in California. *Plant Dis.* 97:259–70
119. Mauch-Mani B, Mauch F. 2005. The role of abscisic acid in plant-pathogen interactions. *Curr. Opin. Plant Biol.* 8:409–14
120. McDonald KL, Cahill DM. 1999. Influence of abscisic acid and the abscisic acid biosynthesis inhibitor, norflurazon, on interactions between *Phytophthora sojae* and soybean (*Glycine max*). *Eur. J. Plant Pathol.* 105:651–58
121. Melotto M, Underwood W, He SY. 2008. Role of stomata in plant innate immunity and foliar bacterial diseases. *Annu. Rev. Phytopathol.* 46:101–22
122. Meng XZ, Xu J, He YX, Yang KY, Mordorski B, et al. 2013. Phosphorylation of an ERF transcription factor by *Arabidopsis* MPK3/MPK6 regulates plant defense gene induction and fungal resistance. *Plant Cell* 25:1126–42

123. Michniewicz M, Czerwinska E, Rozej B. 1990. Interaction of abscisic acid and ethylene in relation to disease development in wheat seedlings infected by *Fusarium culmorum* (W.G.Sm.) Sacc. *Acta Physiol. Plant.* 12:41–48
124. Mittler R. 2006. Abiotic stress, the field environment and stress combination. *Trends Plant Sci.* 11:15–19
125. Mittler R, Vanderauwera S, Gollery M, Van Breusegem F. 2004. Reactive oxygen gene network of plants. *Trends Plant Sci.* 9:490–98
126. Mittler R, Vanderauwera S, Suzuki N, Miller G, Tognetti VB, et al. 2011. ROS signaling: the new wave? *Trends Plant Sci.* 16:300–9
127. Mohr PG, Cahill DM. 2001. Relative roles of glyceollin, lignin and the hypersensitive response and the influence of ABA in compatible and incompatible interactions of soybeans with *Phytophthora sojae*. *Physiol. Mol. Plant Pathol.* 58:31–41
128. Mohr PG, Cahill DM. 2003. Abscisic acid influences the susceptibility of *Arabidopsis thaliana* to *Pseudomonas syringae* pv. *tomato* and *Peronospora parasitica*. *Funct. Plant Biol.* 30:461–69
129. Mohr PG, Cahill DM. 2007. Suppression by ABA of salicylic acid and lignin accumulation and the expression of multiple genes, in *Arabidopsis* infected with *Pseudomonas syringae* pv. *tomato*. *Funct. Integr. Genomics* 7:181–91
130. Montillet J-L, Leonhardt N, Mondy S, Tranchimand S, Rumeau D, et al. 2013. An abscisic acid-independent oxylipin pathway controls stomatal closure and immune defense in *Arabidopsis*. *PLoS Biol.* 11:e1001513
131. Moore GA. 1987. The variation of bark-wood bond strength with moisture content of *Pinus radiata* and three eucalypt species during storage. *Aust. For. Res.* 17:73–78
132. Moore T, Martineau B, Bostock RM, Lincoln JE, Gilchrist DG. 1999. Molecular and genetic characterization of ethylene involvement in mycotoxin-induced plant cell death. *Physiol. Mol. Plant Pathol.* 54:73–85
133. Munns R. 2005. Genes and salt tolerance: bringing them together. *New Phytol.* 167:645–63
134. Munns R, Tester M. 2008. Mechanisms of salinity tolerance. *Annu. Rev. Plant Biol.* 59:651–81
135. Nakashita H, Yasuda M, Nitta T, Asami T, Fujioka S, et al. 2003. Brassinosteroid functions in a broad range of disease resistance in tobacco and rice. *Plant J.* 33:887–98
136. Nambara E, Marion-Poll A. 2005. Abscisic acid biosynthesis and catabolism. *Annu. Rev. Plant Biol.* 56:165–85
137. Navarro L, Bari R, Achard P, Lison P, Nemri A, et al. 2008. DELLAs control plant immune responses by modulating the balance and salicylic acid signaling. *Curr. Biol.* 18:650–55
138. Navarro L, Dunoyer P, Jay F, Arnold B, Dharmasiri N, et al. 2006. A plant miRNA contributes to antibacterial resistance by repressing auxin signaling. *Science* 312:436–39
139. Nguyen VT, Vuong TD, VanToai T, Lee JD, Wu X, et al. 2012. Mapping of quantitative trait loci associated with resistance to *Phytophthora sojae* and flooding tolerance in soybean. *Crop Sci.* 52:2481–93
140. Nishiyama R, Watanabe Y, Fujita Y, Le DT, Kojima M, et al. 2011. Analysis of cytokinin mutants and regulation of cytokinin metabolic genes reveals important regulatory roles of cytokinins in drought, salt and abscisic acid responses, and abscisic acid biosynthesis. *Plant Cell* 23:2169–83
141. Osakabe Y, Arinaga N, Umezawa T, Katsura S, Nagamachi K, et al. 2013. Osmotic stress responses and plant growth controlled by potassium transporters in *Arabidopsis*. *Plant Cell* 25:609–24
142. Osakabe Y, Yamaguchi-Shinozaki K, Shinozaki K, Tran LSP. 2013. Sensing the environment: key roles of membrane-localized kinases in plant perception and response to abiotic stress. *J. Exp. Bot.* 64:445–58
143. Pareek A, Sopory SK, Bohnert HJ, Govindjee X, eds. 2010. *Abiotic Stress Adaptation in Plants: Physiological, Molecular and Genomic Foundation*. Dordrecht, Neth.: Springer
144. Park JE, Park JY, Kim YS, Staswick PE, Jeon J, et al. 2007. GH3-mediated auxin homeostasis links growth regulation with stress adaptation response in *Arabidopsis*. *J. Biol. Chem.* 282:10036–46
145. Park JM, Park C-J, Lee S-B, Ham B-K, Shin R, Paek K-H. 2001. Overexpression of the tobacco *Tsi1* gene encoding an EREBP/AP2-type transcription factor enhances resistance against pathogen attack and osmotic stress in tobacco. *Plant Cell* 13:1035–46
146. Peleg Z, Blumwald E. 2011. Hormone balance and abiotic stress tolerance in crop plants. *Curr. Opin. Plant Biol.* 14:290–95

147. Phillips JR, Dalmay T, Bartels D. 2007. The role of small RNAs in abiotic stress. *FEBS Lett.* 581:3592–97
148. Pieterse CM, Van der Does D, Zamioudis C, Leon-Reyes A, Van Wees SC. 2012. Hormonal modulation of plant immunity. *Annu. Rev. Cell Dev. Biol.* 28:489–521
149. Potters G, Pasternak TP, Guisez Y, Jansen MAK. 2009. Different stresses, similar morphogenic responses: integrating a plethora of pathways. *Plant Cell Environ.* 32:158–69
150. Prasch CM, Sonnewald U. 2013. Simultaneous application of heat, drought, and virus to *Arabidopsis* plants reveals significant shifts in signaling networks. *Plant Physiol.* 162:1849–66
151. Pumplun N, Voinnet O. 2013. RNA silencing suppression by plant pathogens: defence, counter-defence and counter-counter-defence. *Nat. Rev. Microbiol.* 11:745–60
152. Pye MF, Hakuno F, MacDonald JD, Bostock RM. 2013. Induced resistance in tomato by SAR activators during predisposing salinity stress. *Front. Plant Sci.* 4:116
153. Ren B, Liang Y, Deng Y, Chen Q, Zhang J, et al. 2009. Genome-wide comparative analysis of type-A *Arabidopsis* response regulator genes by overexpression studies reveals their diverse roles and regulatory mechanisms in cytokinin signaling. *Cell Res.* 19:1178–90
154. Reyes JL, Chua N-H. 2007. ABA induction of miR159 controls transcript levels of two MYB factors during *Arabidopsis* seed germination. *Plant J.* 49:592–606
155. Rivero RM, Kojima M, Gepstein A, Sakakibara H, Mittler R, et al. 2007. Delayed leaf senescence induces extreme drought tolerance in a flowering plant. *Proc. Natl. Acad. Sci. USA* 104:19631–36
156. Roatti B, Perazzolli M, Gessler C, Pertot I. 2013. Abiotic stresses affect *Trichoderma harzianum* T39-induced resistance to downy mildew in grapevine. *Phytopathology* 103:1227–34
157. Robert-Seilaniantz A, Grant M, Jones JDG. 2011. Hormone crosstalk in plant disease and defense: more than just jasmonate-salicylate antagonism. *Annu. Rev. Phytopathol.* 49:317–43
158. Rock CD, Sakata Y, Quatrano RS. 2010. Stress signaling I: the role of abscisic acid (ABA). See Ref. 143, pp. 33–73.
159. Roubtsova TV, Bostock RM. 2009. Episodic abiotic stress as a potential contributing factor to onset and severity of disease caused by *Phytophthora ramorum* in *Rhododendron* and *Viburnum*. *Plant Dis.* 93:912–18
160. Sagi M, Fluhr R. 2006. Production of reactive oxygen species by plant NADPH oxidases. *Plant Physiol.* 141:336–40
161. Sanchez DH, Pieckenstein FL, Szymanski J, Erban A, Bromke M, et al. 2011. Comparative functional genomics of salt stress in related model and cultivated plants identifies and overcomes limitations to translational genomics. *PLoS ONE* 6:e17094
162. Sandermann H. 1996. Ozone and plant health. *Annu. Rev. Phytopathol.* 34:347–66
163. Savchenko T, Kolla V, Wang C-Q, Nasafi Z, Hicks D, et al. 2014. Functional convergence of oxylipin and ABA pathways controls stomatal closure in response to drought. *Plant Physiol.* 164:1151–60
164. Schenke D, Bottcher C, Scheel D. 2011. Crosstalk between abiotic ultraviolet-B stress and biotic (flg22) stress signalling in *Arabidopsis* prevents flavonol accumulation in favor of pathogen defence compound production. *Plant Cell Environ.* 34:1849–64
165. Schoeneweiss DF. 1975. Predisposition, stress, and plant disease. *Annu. Rev. Phytopathol.* 13:193–211
166. Schoeneweiss DF. 1978. Water stress as a predisposing factor in plant disease. In *Water Deficits and Plant Growth*, ed. TT Kozlowski, pp. 61–99. New York: Academic
167. Schoeneweiss DF. 1981. The role of environmental stress in diseases of woody plants. *Plant Dis.* 65:308–14
168. Shaw RE, Meyer WS, McNeill A, Tyerman SD. 2013. Waterlogging in Australian agricultural landscapes: a review of plant responses and crop models. *Crop Pasture Sci.* 64:549–62
169. Shivaprasad PV, Chen H-M, Patel K, Bond DM, Santos BACM, Baulcombe DC. 2012. A microRNA superfamily regulates nucleotide binding site-leucine-rich repeats and other mRNAs. *Plant Cell* 24:859–74
170. Shkolnik-Inbar D, Bar-Zvi D. 2010. ABI4 mediates abscisic acid and cytokinin inhibition of lateral root formation by reducing polar auxin transport in *Arabidopsis*. *Plant Cell* 22:3560–73
171. Snoeijsers SS, Perez-Garcia A, Joosten M, De Wit P. 2000. The effect of nitrogen on disease development and gene expression in bacterial and fungal plant pathogens. *Eur. J. Plant Pathol.* 106:493–506
172. Sorauer P, Lindau G, Reh L. 1922. *Manual of Plant Diseases*. Wilkes-Barre, PA: Record Press

173. Staswick PE, Tiryaki I, Rowe ML. 2002. Jasmonate response locus *JARI* and several related *Arabidopsis* genes encode enzymes of the firefly luciferase superfamily that show activity on jasmonic, salicylic, and indole-3-acetic acids in an assay for adenylation. *Plant Cell* 14:1405–15
174. Stermer BA, Hammerschmidt R. 1987. Association of heat-shock induced resistance to disease with increased accumulation of insoluble extensin and ethylene synthesis. *Physiol. Mol. Plant Pathol.* 31:453–61
175. Stevens J, Senaratna T, Sivasithamparam K. 2006. Salicylic acid induces salinity tolerance in tomato (*Lycopersicon esculentum* cv. Roma): associated changes in gas exchange, water relations and membrane stabilisation. *Plant Growth Regul.* 49:77–83
176. Stratmann J. 2003. Ultraviolet-B radiation co-opts defense signaling pathways. *Trends Plant Sci.* 8:526–33
177. Sturrock RN, Frankel SJ, Brown AV, Hennon PE, Kliejunas JT, et al. 2011. Climate change and forest diseases. *Plant Pathol.* 60:133–49
178. Sun Y, Li L, Macho AP, Han Z, Hu Z, et al. 2013. Structural basis for flg22-induced activation of the *Arabidopsis* FLS2-BAK1 immune complex. *Science* 342:624–28
179. Sunkar R, Chinnusamy V, Zhu J, Zhu J-K. 2007. Small RNAs as big players in plant abiotic stress responses and nutrient deprivation. *Trends Plant Sci.* 12:301–9
180. Suntio T, Mäkinen K. 2012. Abiotic stress responses promote *Potato virus A* infection in *Nicotiana benthamiana*. *Mol. Plant Pathol.* 13:775–84
181. Taiz L, Zeiger E. 2006. *Plant Physiology*. Sunderland, MA: Sinauer Assoc.
182. Teige M, Scheikl E, Eulgem T, Doczi F, Ichimura K, et al. 2004. The MKK2 pathway mediates cold and salt stress signaling in *Arabidopsis*. *Mol. Cell* 15:141–52
183. Thaler JS, Bostock RM. 2004. Interactions between abscisic-acid-mediated responses and plant resistance to pathogens and insects. *Ecology* 85:48–58
184. Theocharis A, Clement C, Barka EA. 2012. Physiological and molecular changes in plants grown at low temperatures. *Planta* 235:1091–105
185. Ton J, Flors V, Mauch-Mani B. 2009. The multifaceted role of ABA in disease resistance. *Trends Plant Sci.* 14:310–17
186. Truman WM, Bennett MH, Turnbull CGN, Grant MR. 2010. *Arabidopsis* auxin mutants are compromised in systemic acquired resistance and exhibit aberrant accumulation of various indolic compounds. *Plant Physiol.* 152:1562–73
187. Tsutsui T, Kato W, Asada Y, Sako K, Sato T, et al. 2009. DEAR1, a transcriptional repressor of DREB protein that mediates plant defense and freezing stress responses in *Arabidopsis*. *J. Plant Res.* 122:633–43
188. Umezawa T, Hirayama T, Kuromori T, Shinozaki K. 2011. The regulatory networks of plant responses to abscisic acid. In *Plant Responses to Drought and Salinity Stress: Developments in a Post-Genomic Era*, ed. I Turkan, pp. 201–48. New York: Acad. Press
189. Verslues PE, Agarwal M, Katiyar-Agarwal S, Zhu JH, Zhu JK. 2006. Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. *Plant J.* 45:523–39
190. Vicente MRS, Plasencia J. 2011. Salicylic acid beyond defence: its role in plant growth and development. *J. Exp. Bot.* 62:3321–38
191. Vlot AC, Dempsey DMA, Klessig DF. 2009. Salicylic acid, a multifaceted hormone to combat disease. *Annu. Rev. Phytopathol.* 47:177–206
192. Walters DR, McRoberts N. 2006. Plants and biotrophs: a pivotal role for cytokinins? *Trends Plant Sci.* 11:581–86
193. Walters DR, Ratsep J, Havis ND. 2013. Controlling crop diseases using induced resistance: challenges for the future. *J. Exp. Bot.* 64:1263–80
194. Wang D, Pajerowska-Mukhtar K, Culler AH, Dong XN. 2007. Salicylic acid inhibits pathogen growth in plants through repression of the auxin signaling pathway. *Curr. Biol.* 17:1784–90
195. Wang M, Zheng QS, Shen QR, Guo SW. 2013. The critical role of potassium in plant stress response. *Int. J. Mol. Sci.* 14:7370–90
196. Wang Z-Y. 2012. Brassinosteroids modulate plant immunity at multiple levels. *Proc. Natl. Acad. Sci. USA* 109:7–8

197. Wasternack C, Hause B. 2013. Jasmonates: biosynthesis, perception, signal transduction and action in plant stress response, growth and development. An update to the 2007 review in *Annals of Botany*. *Ann. Bot.* 111:1021–58
198. Weyman PD, Pan ZQ, Feng Q, Gilchrist DG, Bostock RM. 2006. A circadian rhythm-regulated tomato gene is induced by arachidonic acid and *Phytophthora infestans* infection. *Plant Physiol.* 140:235–48
199. Wilkinson S, Kudoyarova GR, Veselov DS, Arkhipova TN, Davies WJ. 2012. Plant hormone interactions: innovative targets for crop breeding and management. *J. Exp. Bot.* 63:3499–509
200. Xing Y, Jia W, Zhang J. 2008. AtMKK1 mediates ABA-induced CAT1 expression and H₂O₂ production via AtMPK6-coupled signaling in *Arabidopsis*. *Plant J.* 54:440–51
201. Xu J, Audenaert K, Hofte M, De Vleeschauwer D. 2013. Abscisic acid promotes susceptibility to the rice leaf blight pathogen *Xanthomonas oryzae* pv *oryzae* by suppressing salicylic acid-mediated defenses. *PLoS ONE* 8:e67413
202. Xue-Xuan X, Hong-Bo S, Yuan-Yuan M, Gang X, Jun-Na S, et al. 2010. Biotechnological implications from abscisic acid (ABA) roles in cold stress and leaf senescence as an important signal for improving plant sustainable survival under abiotic-stressed conditions. *Crit. Rev. Biotechnol.* 30:222–30
203. Yamaguchi-Shinozaki K, Shinozaki K. 2006. Transcriptional regulatory networks in cellular responses and tolerance to dehydration and cold stresses. *Annu. Rev. Plant Biol.* 57:781–803
204. Yang D-L, Yao J, Mei C-S, Tong X-H, Zeng L-J, et al. 2012. Plant hormone jasmonate prioritizes defense over growth by interfering with gibberellin signaling cascade. *Proc. Natl. Acad. Sci. USA* 109:E1192–200
205. Yarwood CE. 1959. Predisposition. In *Plant Pathology*, ed. JG Horsfall, AE Dimond, pp. 521–62. New York: Acad. Press
206. Yasuda M, Ishikawa A, Jikumaru Y, Seki M, Umezawa T, et al. 2008. Antagonistic interaction between systemic acquired resistance and the abscisic acid-mediated abiotic stress response in *Arabidopsis*. *Plant Cell* 20:1678–92
207. Yesbergenova Z, Yang G, Oron E, Soffer D, Fluhr R, Sagi M. 2005. The plant Mo-hydroxylases aldehyde oxidase and xanthine dehydrogenase have distinct reactive oxygen species signatures and are induced by drought and abscisic acid. *Plant J.* 42:862–76
208. Yoshioka K, Shinozaki K. 2009. *Signal Crosstalk in Plant Stress Responses*. Ames, IA: Wiley-Blackwell
209. Zhang S, Cai Z, Wang X. 2009. The primary signaling outputs of brassinosteroids are regulated by abscisic acid signaling. *Proc. Natl. Acad. Sci. USA* 106:4543–48
210. Zhao JQ, Li S, Jiang TF, Liu Z, Zhang WW, et al. 2012. Chilling stress: the key predisposing factor for causing *Alternaria alternata* infection and leading to cotton (*Gossypium hirsutum* L.) leaf senescence. *PLoS ONE* 7:e36126
211. Zhu Q-H, Fan L, Liu Y, Xu H, Llewellyn D, Wilson I. 2013. miR482 regulation of NBS-LRR defense genes during fungal pathogen infection in cotton. *PLoS ONE* 8:e84390
212. Zhu Y, Qian W, Hua J. 2010. Temperature modulates plant defense responses through NB-LRR proteins. *PLoS Pathog.* 6:e1000844



Contents

How Way Leads on to Way <i>Isaac Barash</i>	1
Harnessing Population Genomics to Understand How Bacterial Pathogens Emerge, Adapt to Crop Hosts, and Disseminate <i>Boris A. Vinatzer, Caroline L. Monteil, and Christopher R. Clarke</i>	19
New Insights into Mycoviruses and Exploration for the Biological Control of Crop Fungal Diseases <i>Jiatao Xie and Daohong Jiang</i>	45
Altering the Cell Wall and Its Impact on Plant Disease: From Forage to Bioenergy <i>Qiao Zhao and Richard A. Dixon</i>	69
Network Modeling to Understand Plant Immunity <i>Oliver Windram, Christopher A. Penfold, and Katherine J. Denby</i>	93
The Role of Trees in Agroecology and Sustainable Agriculture in the Tropics <i>Roger R.B. Leakey</i>	113
Plant-Parasitic Nematode Infections in Rice: Molecular and Cellular Insights <i>Tina Kyndt, Diana Fernandez, and Godelieve Gheysen</i>	135
Mechanisms of Nutrient Acquisition and Utilization During Fungal Infections of Leaves <i>Jessie Fernandez, Margarita Marroquin-Guzman, and Richard A. Wilson</i>	155
Governing Principles Can Guide Fungicide-Resistance Management Tactics <i>Frank van den Bosch, Richard Oliver, Femke van den Berg, and Neil Paveley</i>	175
Virus Infection Cycle Events Coupled to RNA Replication <i>Pooja Saxena and George P. Lomonossoff</i>	197

Novel Insights into Rice Innate Immunity Against Bacterial and Fungal Pathogens <i>Wende Liu, Jinling Liu, Lindsay Triplett, Jan E. Leach, and Guo-Liang Wang</i>	213
The Activation and Suppression of Plant Innate Immunity by Parasitic Nematodes <i>Aska Goverse and Geert Smant</i>	243
Protein Kinases in Plant-Pathogenic Fungi: Conserved Regulators of Infection <i>David Turrà, David Segorbe, and Antonio Di Pietro</i>	267
Speciation in Fungal and Oomycete Plant Pathogens <i>Silvia Restrepo, Javier F. Tabima, Maria F. Mideros, Niklaus J. Grünwald, and Daniel R. Matute</i>	289
The ABCs and 123s of Bacterial Secretion Systems in Plant Pathogenesis <i>Jeff H. Chang, Darrell Desveaux, and Allison L. Creason</i>	317
Induced Systemic Resistance by Beneficial Microbes <i>Corné M.J. Pieterse, Christos Zamioudis, Roeland L. Berendsen, David M. Weller, Saskia C.M. Van Wees, and Peter A.H.M. Bakker</i>	347
Fifty Years Since <i>Silent Spring</i> <i>Lynn Epstein</i>	377
Localizing Viruses in Their Insect Vectors <i>Stéphane Blanc, Martin Drucker, and Marilyne Uzest</i>	403
Plant Cell Wall-Degrading Enzymes and Their Secretion in Plant-Pathogenic Fungi <i>Christian P. Kubicek, Trevor L. Starr, and N. Louise Glass</i>	427
Meta-Analysis and Other Approaches for Synthesizing Structured and Unstructured Data in Plant Pathology <i>H. Scherm, C.S. Thomas, K.A. Garrett, and J.M. Olsen</i>	453
Networks and Plant Disease Management: Concepts and Applications <i>M.W. Shaw and M. Pautasso</i>	477
Small RNAs: A New Paradigm in Plant-Microbe Interactions <i>Arne Weiberg, Ming Wang, Marschal Bellinger, and Hailing Jin</i>	495
Predisposition in Plant Disease: Exploiting the Nexus in Abiotic and Biotic Stress Perception and Response <i>Richard M. Bostock, Matthew F. Pye, and Tatiana V. Roubtsova</i>	517

Susceptibility Genes 101: How to Be a Good Host <i>Chris C.N. van Schie and Frank L.W. Takken</i>	551
Horizontal Gene Transfer in Eukaryotic Plant Pathogens <i>Darren Soanes and Thomas A. Richards</i>	583

Errata

An online log of corrections to *Annual Review of Phytopathology* articles may be found at <http://www.annualreviews.org/errata/phyto>



ANNUAL REVIEWS

It's about time. Your time. It's time well spent.

New From Annual Reviews:

Annual Review of Statistics and Its Application

Volume 1 • Online January 2014 • <http://statistics.annualreviews.org>

Editor: **Stephen E. Fienberg**, *Carnegie Mellon University*

Associate Editors: **Nancy Reid**, *University of Toronto*

Stephen M. Stigler, *University of Chicago*

The *Annual Review of Statistics and Its Application* aims to inform statisticians and quantitative methodologists, as well as all scientists and users of statistics about major methodological advances and the computational tools that allow for their implementation. It will include developments in the field of statistics, including theoretical statistical underpinnings of new methodology, as well as developments in specific application domains such as biostatistics and bioinformatics, economics, machine learning, psychology, sociology, and aspects of the physical sciences.

Complimentary online access to the first volume will be available until January 2015.

TABLE OF CONTENTS:

- *What Is Statistics?* Stephen E. Fienberg
- *A Systematic Statistical Approach to Evaluating Evidence from Observational Studies*, David Madigan, Paul E. Stang, Jesse A. Berlin, Martijn Schuemie, J. Marc Overhage, Marc A. Suchard, Bill Dumouchel, Abraham G. Hartzema, Patrick B. Ryan
- *The Role of Statistics in the Discovery of a Higgs Boson*, David A. van Dyk
- *Brain Imaging Analysis*, F. DuBois Bowman
- *Statistics and Climate*, Peter Guttorp
- *Climate Simulators and Climate Projections*, Jonathan Rougier, Michael Goldstein
- *Probabilistic Forecasting*, Tilmann Gneiting, Matthias Katzfuss
- *Bayesian Computational Tools*, Christian P. Robert
- *Bayesian Computation Via Markov Chain Monte Carlo*, Radu V. Craiu, Jeffrey S. Rosenthal
- *Build, Compute, Critique, Repeat: Data Analysis with Latent Variable Models*, David M. Blei
- *Structured Regularizers for High-Dimensional Problems: Statistical and Computational Issues*, Martin J. Wainwright
- *High-Dimensional Statistics with a View Toward Applications in Biology*, Peter Bühlmann, Markus Kalisch, Lukas Meier
- *Next-Generation Statistical Genetics: Modeling, Penalization, and Optimization in High-Dimensional Data*, Kenneth Lange, Jeanette C. Papp, Janet S. Sinsheimer, Eric M. Sobel
- *Breaking Bad: Two Decades of Life-Course Data Analysis in Criminology, Developmental Psychology, and Beyond*, Elena A. Erosheva, Ross L. Matsueda, Donatello Telesca
- *Event History Analysis*, Niels Keiding
- *Statistical Evaluation of Forensic DNA Profile Evidence*, Christopher D. Steele, David J. Balding
- *Using League Table Rankings in Public Policy Formation: Statistical Issues*, Harvey Goldstein
- *Statistical Ecology*, Ruth King
- *Estimating the Number of Species in Microbial Diversity Studies*, John Bunge, Amy Willis, Fiona Walsh
- *Dynamic Treatment Regimes*, Bibhas Chakraborty, Susan A. Murphy
- *Statistics and Related Topics in Single-Molecule Biophysics*, Hong Qian, S.C. Kou
- *Statistics and Quantitative Risk Management for Banking and Insurance*, Paul Embrechts, Marius Hofert

Access this and all other Annual Reviews journals via your institution at www.annualreviews.org.

ANNUAL REVIEWS | Connect With Our Experts

Tel: 800.523.8635 (US/CAN) | Tel: 650.493.4400 | Fax: 650.424.0910 | Email: service@annualreviews.org

